

AD-A096 970

AEROSPACE CORP EL SEGUNDO CA SPACE SCIENCES LAB  
ENERGETIC-PARTICLE POPULATIONS AND COSMIC-RAY ENTRY.(U)  
MAR 81 M SCHULZ  
TR-0081(6960-05)-7

F/6 4/1

F04701-80-C-0081


UNCLASSIFIED

SD-TR-81-13

NL

[ CR ]

REV  
05/89/70



END

DATE

FILED

DTIC

LEVEL II

(12)

AD A000000

# Energetic-Particle Populations and Cosmic-Ray Entry

M. SCHULZ  
Space Sciences Laboratory  
Laboratory Operations  
The Aerospace Corporation  
El Segundo, Calif. 90245

DTIC  
ELECTE  
MAR 27 1981  
S D E

17 March 1981

Interim Report

APPROVED FOR PUBLIC RELEASE;  
DISTRIBUTION UNLIMITED

DTIC FILE COPY


Prepared for  
SPACE DIVISION  
AIR FORCE SYSTEMS COMMAND  
Los Angeles Air Force Station  
P.O. Box 92960, Worldway Postal Center  
Los Angeles, Calif. 90009


81 3 27 008

This interim report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract No. F04701-80-C-0081 with the Space Division, P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by G.A. Paulikas, Director, Space Sciences Laboratory. Major George A. Kuck, YLXT, was the project officer for the Mission Oriented Investigation and Experimentation Program.


This report has been reviewed by the Public Affairs Office (PAS) and is releasable to the National Technical Information Service (NITS). At NITS it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

  
George A. Kuck, Major, USAF  
Project Officer

  
Florian P. Meinhardt, Lt Col, USAF  
Director of Advanced Space Development

FOR THE COMMANDER

  
William Goldberg, Colonel, USAF  
Deputy for Technology

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 18 SD-TR-81-13	2. GOVT ACCESSION NO. AD-A096	3. RECIPIENT'S CATALOG NUMBER 970
4. TITLE (and Subtitle) ENERGETIC-PARTICLE POPULATIONS AND COSMIC-RAY ENTRY		5. TYPE OF REPORT & PERIOD COVERED 9 INTERIM technical rept.
7. AUTHOR(s) 10 Michael/Schulz	14 TR-0081(6960-05)-7	6. PERFORMING ORG. REPORT NUMBER 15 F04701-80-C-0081
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Aerospace Corporation El Segundo, Calif. 90245		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Space Division Air Force Systems Command Los Angeles, Calif. 90009		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 11 17 Mar 81 12 1
		13. NUMBER OF PAGES 75
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Cosmic Rays Magnetosphere Radiation Belts		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  This report constitutes a review of recent progress in the study of charged-particle access, transport, and loss with respect to the earth's magnetosphere. A major objective of studying particle access is to identify the sources of trapped and quasi-trapped radiation. Important progress in this area involves the measurement of comparative ionic abundances in the radiation belts. The historic problem of accounting for cosmic-ray cut-off latitudes as a function of particle energy is being solved by the adoption of increasingly		

00 FORM 1473  
(FACSIMILE)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

19. KEY WORDS (Continued)

20. ABSTRACT (Continued)

realistic and accurate models of the magnetospheric B field. Progress in the estimation of particle-transport coefficients (mainly diffusion coefficients) involves the measurement of fluctuating electric and magnetic fields on the ground, at balloon altitudes, and in space. The importance of particle interactions with discrete waveforms (as distinguished from broad-banded spectral noise) is increasingly being recognized. For example, the unsteady magnetospheric convection associated with substorms contributes importantly to radial diffusion, whereas cyclotron resonance with chorus elements and other discrete excitations may contribute importantly to pitch-angle diffusion and (thus) to the loss of energetic particles from the magnetosphere. The role of man-made signals such as radio transmissions and power-line harmonics in this latter process remains uncertain and continues to be debated.

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## PREFACE

This report is based on a reporter-review (that of IAGA Subdivision III-4) that was presented 5 December 1979 at the XVII General Assembly of the International Union of Geodesy and Geophysics (IUGG) in Canberra, Australia. The author is pleased to thank Dr. C.-G. Fälthammar (chairman of IAGA Division III) and Dr. L. J. Lanzerotti (the elected reporter-reviewer for Topic III-4) for the invitation to present this review. He is also pleased to thank Dr. G. Rostoker and Dr. A. Nishida for suggesting that the written report be published in this journal. The author is pleased to thank the American Geophysical Union for partially subsidizing the trip to Australia under a group travel grant from the U. S. National Science Foundation. Finally, it is a pleasure to thank Inge Scanlan and Lilia Francewar for typing the final manuscript and supplementary bibliography.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
<b>A</b>	

## CONTENTS

PREFACE.....	1
INTRODUCTION.....	5
ENERGETIC-PARTICLE ACCESS OF THE MAGNETOSPHERE.....	7
ENERGETIC-PARTICLE TRANSPORT IN THE MAGNETOSPHERE.....	19
ENERGETIC-PARTICLE LOSS PROCESSES.....	23
KINEMATICAL EFFECTS.....	35
EPILOGUE.....	37
REFERENCES.....	39

## FIGURES

1.	Three-dimensional "L-mode" Mass Spectra Taken Near Noon on 29 July 1977.....	9
2.	Radial Profile of Plasma Composition (0.9-13.9 keV/Charge) During Magnetic Storm of 29 July 1977 (Geiss <u>et al.</u> , 1978).....	10
3.	Relative Probability-of-Occurrence of an Upward-Flowing Ion Beam as a Function of Magnetic Local Time in the Altitude Range 6000-8000 km, Normalized to 1.0 for the Interval 15-18 hr MLT (Ghielmetti <u>et al.</u> , 1978).....	11
4.	Schematic Illustration of Electrostatic Equipotentials In and Near an Auroral Arc (Swift <u>et al.</u> , 1976).....	12
5.	Contours of Constant Phase-Space Density for Auroral Electrons, 12 August 1976 (UT = 12120.2 $\pm$ 9.0 sec), as Determined by Croley <u>et al.</u> (1978).....	14
6.	Representative Trajectories of a 5-MeV Proton, as Computed by Blake and Friesen (1979) and Projected Onto the Equatorial Plane of a Model B Field.....	15
7.	Contours of Constant Phase-Space Density at L = 4 (Dusk Meridian) for an Isotropic Source of Temperature 2 keV Placed at L = 10 in the Presence of a 90-kV Cross-Magnetospheric Potential Drop Distributed as $L^2 \sin \phi$ , Where $\phi$ is the Local Time.....	24
8.	Equatorial Growth Rates for Ion-Cyclotron Waves (Electro- Magnetic) at L = 4 (Dusk Meridian), as Computed by Kaye <u>et al.</u> (1979) for Conditions Described in Figure 7 and its Caption, But With Various Source Temperatures.....	25
9.	Average Electron Flux in the Global Loss Cone During the Year 1969 at Energies 139-312 keV, L = 1.5-4.0 (Vampola, 1977).....	27
10.	Mapping of Local Truncation in Pitch-Angle Distribution to 100-km Loss Cone at Smaller Longitude (Vampola and Kuck, 1978).....	28
11.	Dependence of Equatorial Pitch Angle, Corresponding to Local Truncation of Pitch-Angle Distribution, Upon L in the Energy Channel on Which the Effect is Centered (Vampola and Kuck, 1978).....	29
12.	Total Change in Equatorial Pitch Angle Accumulated by Electrons of Various Initial Phases $\phi_0$ After Interaction With Whistler-Mode Wave of Specified Amplitude at L = 4 (Inan <u>et al.</u> , 1978).....	32



## INTRODUCTION

The purpose of this work is to review recent theoretical and observational developments in the study of energetic charged-particle populations in the earth's magnetosphere. More specifically, the intended focus of this review is on developments that have occurred since the Third General Scientific Assembly of IAGA, held in Seattle toward the end of August 1977. This is not the sort of review in which the author seeks to identify the major problems that have been solved during the past few years and to specify the outstanding new problems that have emerged during the same period and now require solution. Progress in magnetospheric physics does not occur in accordance with such a stereotype. It seems instead that the basic questions addressed by magnetospheric physicists remain the same from decade to decade. What evolves over the years is the sophistication with which the basic questions are formulated, and also the detail in which answers are sought. This situation is a profound tribute to the pioneers of space research for their ability to identify the outstanding problems of magnetospheric physics at an early date. The evolving standard of sophistication with which answers to these basic questions are sought and delivered is a continuing tribute to the space-physics community as a whole.

The basic questions addressed in radiation-belt physics involve the access, transport, and loss of the charged particles with respect to the earth's magnetosphere. The question of access includes the problem of identifying particle sources. The question of transport includes the problem of identifying both diffusive and non-diffusive mechanisms by which one or more of the adiabatic invariants of a charged particle can change value. The question of particle loss includes the problem of

identifying the mechanisms of pitch-angle diffusion, the size of the loss cone, the rate of energy degradation by various processes, and the lifetime against charge exchange. It is clear from this description that the questions of particle access, transport, and loss overlap somewhat in scope. There is no harm in this, since the objective of radiation-belt physics is to provide a coherent, comprehensive, and quantitative description of the energetic particle populations in the earth's magnetosphere. However, the allocation of certain topics to the "access" and "loss" categories (rather than to "transport") in the present work is necessarily arbitrary in some cases.

## ENERGETIC-PARTICLE ACCESS TO THE MAGNETOSPHERE

It has long been realized that a measurement of comparative ionic abundances in the magnetosphere carries implications concerning the source of the particles observed. The basic concept is that various conceivable sources for geomagnetically trapped radiation have different ionic-abundance ratios, and that a ratio at the source should be reflected in the magnetospheric abundance ratio under certain conditions. The situation is somewhat complicated by the possibility of differences in transport rates within the magnetosphere, but there is considerable truth in the basic concept. Cornwall and Schulz (1979) have recently reviewed this concept and its limitations, together with some of the early observations of the He/H ionic abundance ratios in the magnetosphere and the puzzles that these observations presented. Spjeldvik (1979) has recently reviewed the subject from a more theoretical viewpoint, with emphasis on comparative charge-state abundances of heavier ions such as oxygen. <sup>Prangé (1978),</sup> Johnson (1979), Shelley (1979), and Hultqvist (1979) have reviewed past observations of energetic ions at altitudes  $\leq 10^4$  km in the auroral regions and have thus emphasized ions of ionospheric origin. Young (1979) has provided a review of these, as well as more recent observations, and has thus approached the subject from a magnetospheric perspective.

Perhaps the major observational development in this area of magnetospheric research in recent years has been the placement of the ISEE and GEOS spacecraft in near-equatorial orbits that extend to high L values. Preliminary results from the instruments onboard GEOS-1, ISEE-1, and ISEE-2 appeared in six special issues of Space Science Reviews during 1978-79. GEOS-2 and ISEE-3 were launched following the Thirteenth ESLAB Symposium (held 5-7 June 1978 at Innsbruck, Austria), at which the

preliminary results from the three earlier spacecraft were first publicly presented. The major results on ionic composition were the GEOS results of Geiss et al. (1978). Some of these are illustrated in Figures 1-2. The reader will note that the energies represented here are quite low by radiation-belt standards, and even exclude a major part of the ring-current spectrum. This difficulty arises from an instrument gap that so far has precluded observations from  $\sim 20$  keV/charge to  $\sim 100$  keV/nucleon (e.g., Cornwall and Schulz, 1979). However, the observations of Geiss et al. (1978) at least suggest that oxygen ( $O^+$ ) is an important constituent of the ring current, along with  $H^+$  and (to a lesser extent) helium. These results confirm the observations that Johnson et al. (1977) had made on S3-3 at locations far from the magnetic equator.

It is presumed that energetic ions enter the magnetosphere either by crossing the magnetopause or by following the electric field of an auroral arc. In support of the latter process, Ghielmetti et al. (1978) have reported the common occurrence of upward flowing energetic ions (both  $H^+$  and  $O^+$ ) of ionospheric origin, especially in the afternoon and evening sectors (see Figure 3). The local-time signature of this phenomenon is consistent with the idea (e.g., Lyons, 1980) that auroral electric fields are generated by an inherent "discontinuity" in the magnetospheric convection field. As is illustrated in Figure 4 (Swift et al., 1976), the parallel (to  $B$ ) component of  $E$  tends to point upward (downward) at places where the normal (to  $B$ ) component of  $E$  points toward (away from) the auroral arc. In the case of a dawn-to-dusk electric field, the parallel (to  $B$ ) component of  $E$  must therefore point upward in the PM sector and downward in the AM sector, as the results shown in Figure 3 suggest.

The signature of an auroral electric field with a strong upward component parallel to  $B$  is evident in the high-resolution velocity-space distributions of ions and electrons compiled by Croley et al. (1978). The results, presented in the form of contour plots on

GEOS/ICE  
 DAY 210  
 29 JULY 1977  
 $L_d = 6.12$   
 $\lambda_d = 3.37$   
 LT = 12.42  
 $K_p = 5^+$




Figure 1. Three-dimensional "L-mode" mass spectra taken near noon on 29 July 1977. This was during the main phase of a magnetic storm for which  $D_{st}$  reached  $-100 \gamma$  (at 0700 UT). Abscissa = mass/charge; ordinate = differential flux per unit charge (Geiss et al., 1973).




Figure 2. Radial profile of plasma composition (0.9-13.9 keV/charge) during magnetic storm of 29 July 1977 (Geiss et al., 1978).




Figure 3. Relative probability-of-occurrence of an upward-flowing ion beam as a function of magnetic local time in the altitude range 6000-8000 km, normalized to 1.0 for the interval 15-18 hr MLT (Ghielmetti et al., 1978).




Figure 4. Schematic illustration of electrostatic equipotentials in and near an auroral arc (Swift et al., 1976). Since  $\underline{E}$  is perpendicular to the equipotential surfaces, an upward  $\underline{E}$  at the center of the arc corresponds to a magnetospheric  $\underline{E}$  that points toward the arc from both sides.




the  $(v_{\parallel}, v_{\perp})$  plane, reveal clearly the boundaries in velocity space that separate adiabatically distinct classes of particles constrained to move along the same field line. This is illustrated for electrons in Figure 5. Data are shown from nearly a full spin of the spacecraft (S3-3) as a precaution against aliasing; to the extent that the distributions in the upper and lower half-planes agree, the distribution thus determined is presumably a good (i.e., instantaneous) one. Torbert and Mozer (1978) observed strong localized electric fields from the same spacecraft in association with particle signatures (Mizera and Fennell, 1977; 1978) of this type.

For ions and electrons that originate not in the ionosphere but in interplanetary space, the mode of access involves crossing the magnetopause somewhere. For particles of moderate energy ( $\leq 10$  keV) this can be a very complicated process indeed. Such particles might enter the boundary-layer convection pattern at the polar cleft or elsewhere on the magnetopause, become part of the plasma mantle, and thence be convected into the plasma sheet. Once in the plasma sheet, such particles might well be energized by both adiabatic and non-adiabatic processes during convection and diffusion into the region of geomagnetically trapped radiation.

Particle access through the magnetopause at substantially higher energies is facilitated by intrinsic violation of the first adiabatic invariant. This occurs wherever the scale size of inhomogeneities in  $\underline{B}$  is smaller than the gyro-radius of the particle. While the magnetopause itself represents such an inhomogeneity (of ideally vanishing scale size), the probability that the magnetosphere will actually trap an incident particle can be enhanced by the additional presence of interior inhomogeneities in  $\underline{B}$ . This is illustrated in Figure 6, in which particle trajectories (as projected onto the equatorial plane of a model  $\underline{B}$  field) are modified by encounters with a field-aligned wedge of drastically reduced  $|\underline{B}|$ . The work illustrated is that of Blake and Friesen (1979). The presence of such diamagnetic "blobs" in the magnetosphere could scatter an incident

ELECTRONS  
DAY 225

UT 12111.20 - 12129.20



1	.100E-04
2	.215E-04
3	.484E-04
4	.100E-03
5	.215E-03
6	.464E-03
7	.100E-02
8	.215E-02
9	.484E-02
0	.100E-01
A	.215E-01
B	.484E-01
C	.100E+00
D	.215E+00
E	.484E+00
F	.100E+01
G	.215E+01
H	.484E+01
I	.100E+02
J	.215E+02

Figure 5. Contours of constant phase-space density for auroral electrons. 12 August 1976 (UT = 12120.2 $\pm$ 9.0 sec), as determined by Croley et al. (1978). Hyperbola and ellipse demarcate regions of velocity space adiabatically accessible to the foot of the field line (100-km altitude) and the equator, respectively, in the presence of the parallel (to B) electric field that has been inferred from the S3-3 data for this time interval.




Figure 6. Representative trajectories of a 5-MeV proton, as computed by Blake and Friesen (1979) and projected onto the equatorial plane of a model B field. The shaded region corresponds to a diamagnetic wedge in which the magnitude of B has been reduced by a factor of ten.

particle's guiding center either toward or away from the magnetopause with roughly equal probability. However, since the source of "cosmic" rays is external to the magnetosphere, the net effect of such scattering is to facilitate particle access on the average. Many of the incident "cosmic" rays are likely to be relativistic electrons emitted from Jupiter's magnetosphere (Baker et al., 1979).

Traditional computations of cosmic-ray cutoffs necessarily omit unpredictable features such as diamagnetic "blobs" from the field model. However, such computations commonly failed (by several degrees of latitude) to account quantitatively for the observed cutoffs because the B-field models previously in use were not accurate enough. Puzzled investigators desperately postulated arbitrary diffusion models in order to account for the discrepancies. However, recent work by Pfitzer (1979) has shown that an improved B-field model is all that was ever required. The field model that he successfully used for computing cosmic-ray cutoffs, such as those illustrated in Table 1, is an unpublished improvement on the model of Olson and Pfitzer (1974). The new field model permits a realistic angle ( $\zeta \approx 90^\circ$ ) between the solar-wind velocity and the dipole axis, and (more importantly) provides a better global fit to the observed B field. Ring currents, tail currents, and magnetopause currents are included as before, but are modeled more accurately.

Debrunner et al. (1979) successfully utilized cosmic-ray cutoff calculations in conjunction with neutron-monitor data to determine the ring-current radius during a magnetic storm. Darchieva et al. (1978) had previously made an observational study of the variation of cutoff latitude with ring-current intensity during major geomagnetic disturbances. Other important trajectory studies include purely theoretical works involving neutral points (Amano and Tsuda, 1978) and current sheets (Cowley, 1978). Gall and Bravo (1979) have made significant studies of solar-proton trajectories in the geo-

Table 1. Observed and computed cutoff latitudes  
for 5-MeV protons in various field models,  
as summarized by Pfitzer (1979).

Model (Reference)	Midnight	Dawn	Noon	Dusk
Gall <u>et al.</u> (1968)	68°		72°	
Olson (1970)	69°		74°	
Olson-Pfitzer (1974)	66°	70°	69°	71°
Pfitzer (1979), $\zeta = 55^\circ$	65°	67°	68°	67°
Pfitzer (1979), $\zeta = 90^\circ$	<u>66°</u>	<u>67°</u>	<u>70°</u>	<u>67°</u>
Pfitzer (1979), $\zeta = 125^\circ$	66°	67°	68°	67°
Observations, extrapolated to $\zeta = 90^\circ$ , $K_p = 0$	66.4° ±0.4°	67.5° ±0.9°	70.4° ±0.6°	67.5° ±0.6°

magnetic tail. Detailed trajectory studies of charged-particle motion near the magnetopause would be quite welcome, now that one has access to three-dimensional velocity-space distributions (Frank et al., 1978; Williams et al., 1978) from GEOS and ISEE.

The foregoing processes entail no change of particle identity. However, Blake and Friesen (1977) have described a mechanism whereby an incident heavy ion ( $O^+$ , for example) is stripped of one or more of its remaining electrons as the incident ion grazes the atmosphere near a magnetic mirror point somewhere above the cutoff latitude for particles of that rigidity. By virtue of becoming multiply charged, the ion thus has its rigidity greatly reduced and becomes trapped in the geomagnetic field. There is even a good chance that the ion subsequently has a small enough gyro-radius to obey the laws of adiabatic charged-particle motion.

## ENERGETIC-PARTICLE TRANSPORT IN THE MAGNETOSPHERE

The technical definition of transport is the violation of one or more of the adiabatic invariants of charged-particle motion. Transport can be either diffusive or frictional in character and is described by means of the Fokker-Planck equation. One-shot processes such as charge exchange or the beta decay of a neutron are handled by adding source and loss terms to the Fokker-Planck equation for a species. Other processes that may change the particle flux at a given energy, but which conserve all three adiabatic invariants of charged-particle motion, are kinematical in character and are treated as such in the present work.

Lyons and Williams (1980) have recently found that one can account for the storm-time ring current by "transporting" the quiet-time ring current sufficiently inward in L. No additional source of charged particles or energy would be required, as long as the "transport" process conserves the first two adiabatic invariants. The most likely transport mechanism for this, consistent with the laws of physics, is unsteady magnetospheric convection. The consequences of such transport, even if accomplished by a single impulsive enhancement of the convection electric field, are qualitatively similar to the consequences of enhanced radial diffusion.

For particles of much higher energy than the ring current, it is important to consider also the effects of magnetic impulses. As commonly modeled, the effects of magnetic and electric impulses scale differently with particle energy, charge state, and equatorial pitch angle (e.g., Cornwall and Schulz, 1979). In pursuit of this idea, Lanzerotti et al. (1978) have analyzed the spectrum of (presumably) large-scale geomagnetic noise at synchronous altitude (ATS-6). By evaluating the spectral density at

particle drift frequencies, they have estimated the corresponding radial-diffusion coefficient there. The results of Lanzerotti et al. (1978) are model-dependent, of course, but this is a common hazard in space research. It is better to obtain a model-dependent result than none at all, provided that the model has a rational basis. Other predictions of such a model can subsequently be tested against a more comprehensive set of observational data.

Using a similar philosophy, Holzworth and Mozer (1979) have deduced a radial-diffusion coefficient from electric-field fluctuations measured on simultaneous balloon flights at six auroral-zone locations spanning  $180^\circ$  of longitude. In this case the major model-dependent assumption was that B-field lines are equipotentials of the electric field. Since this assumption is likely to fail in an auroral arc, it would perhaps have been better to fly the balloons at a lower magnetic latitude. However, measurements of this type are highly worthwhile, and the work of Holzworth and Mozer (1979) represents an important first step in the field.

Radial diffusion caused by magnetic or electric impulses is known to leave an additional signature on the affected charged-particle populations. Particles dispersed in L by an individual impulse are consequently organized in drift phase by virtue of Liouville's theorem, and this organization manifests itself in the form of drift-periodic echoes in the particle intensities. Chanteur et al. (1977, 1978) have made a systematic study of this effect and have confirmed its observational significance beyond any reasonable doubt. Belian et al. (1978) suggest that proton drift echoes may alternatively result from an "injection" event, by which they presumably mean either a new supply of particles admitted to the magnetosphere or a dawn-to-dusk electrostatic impulse that is confined to the night side of the magnetosphere. Such a possibility is difficult to deny, but it is not supported by the electron-echo observations of Chanteur et al. (1977, 1978).



The notion of "transport" should also include charge-state transitions caused by charge exchange between energetic ions and the earth's atmosphere. This process is described by coupling the Fokker-Planck equations that describe the evolution of phase-space densities for the various charge states of (for example) oxygen. Spjeldvik and Fritz (1978a) have carried out such calculations in order to interpret observational data (Spjeldvik and Fritz, 1978b) on heavy ions in the magnetosphere.

Charge exchange inherently displaces the guiding center of a geomagnetically trapped particle and thus also contributes to a novel form of radial diffusion. An extreme example of such transport is that which enables ring-current protons to traverse the whole trapping region as neutrals and appear again as ions at low altitudes on the magnetic equator. By alternately losing and regaining its charge in the dense atmosphere, a hydrogen ion/atom would thus experience a truly diffusive transport. For heavier ions the transitions among charge states other than zero lead to truly diffusive transport even at ring-current altitudes. The element of randomness is supplied by the gyrophase at which the charge exchange happens to occur, since this gyrophase determines the direction in which the guiding center is displaced.

Certain aspects of energetic-particle transport are complicated enough that numerical computation of the particle trajectories is required. For example, Smith et al. (1979) have numerically traced the drift of particles in the presence of a time-varying convection electric field, and Harel et al. (1979) have numerically computed the evolution of bulk properties of the hot plasma (self-consistently) in a model of magnetospheric substorms. This type of work has a bright future.

Pitch-angle diffusion is a transport process of direct relevance to particle loss from the magnetosphere. It is for this reason that recent developments in the field of pitch-angle diffusion are discussed in the next section rather than here.

## ENERGETIC-PARTICLE LOSS PROCESSES

A major loss mechanism for energetic particles in the earth's magnetosphere is pitch-angle diffusion into the loss cone. This occurs typically by virtue of cyclotron resonance between particles and a broad-banded spectrum of magnetospheric waves. The waves may be either electrostatic or electromagnetic, but the former tend to predominate outside the plasmasphere and the latter inside.

Waves in the magnetospheric plasma commonly originate from instabilities. Kaye et al. (1979) have considered the electromagnetic ion-cyclotron instability of ring-current protons subject to gradient-curvature drift and magnetospheric convection simultaneously. The instability is driven by pitch-angle anisotropy, as is well known. However, Kaye et al. (1979) have found that peculiar features of the growth-rate spectrum (in accord with observation) can be traced to specific cutoffs that are expected to appear in the phase-space density (see Figure 7) as the ring-current protons drift inward by convection from a source at  $L = 10$ . Contours of constant normalized growth rate  $\gamma/\Omega_p$  at  $L = 4$  are shown in Figure 8 for various source temperatures ( $\gamma$  = temporal growth rate,  $\Omega_p/2\pi$  = proton gyrofrequency,  $\omega/2\pi$  = wave frequency). It is significant, for example, that increasing the source temperature does not necessarily increase the growth rate. Only a small minority of the protons in a 4-keV Maxwellian are able to drift from  $L = 10$  to  $L = 4$  while conserving the adiabatic invariants of charged-particle motion.

Waves generated by instabilities such as that studied by Kaye et al. (1979) are expected to produce pitch-angle diffusion, which (in the presence of an atmospheric loss cone) results in local precipitation. The precipitation in this case occurs mainly in the




Figure 7. Contours of constant phase-space density at  $L = 4$  (dusk meridian) for an isotropic source of temperature 2 keV placed at  $L = 10$  in the presence of a 90-kV cross-magnetospheric potential drop distributed as  $L^2 \sin \varphi$ , where  $\varphi$  is the local time. The particles are protons, and the axes (as well as the contours) have been normalized arbitrarily (Kaye *et al.*, 1979).




Figure 8. Equatorial growth rates for ion-cyclotron waves (electromagnetic) at  $L = 4$  (dusk meridian), as computed by Kaye et al. (1979) for conditions described in Figure 7 and its caption, but with various source temperatures. Wave frequencies and growth rates are normalized with respect to the proton gyrofrequency, as indicated.

dusk sector, i.e., is organized with respect to local time. Vampola (1977) has found, on the other hand, that slot-region electron precipitation is organized with respect to geographic longitude (see Figure 9) if one averages over local time. Vampola (1977) suggests that precipitation maxima are associated with (and thus caused by) major VLF transmitters. It is difficult to test this idea statistically from the data in Figure 9, and it would have been better to use an earth-anchored magnetic longitude for organizing the data. However, the idea that anthropogenic processes actually lead to significant particle precipitation is indeed a provocative one.

A cleaner but more subtle example of transmitter-induced pitch-angle diffusion is shown in Figure 10 (Vampola and Kuck, 1978). In this case the authors observe a pitch-angle distribution that is truncated well outside the local loss cone at east (geographic) longitudes  $> 100^\circ$ . However, the equatorial pitch angle of the truncation consistently corresponds to the loss cone that would have prevailed (assuming precipitation at 100-km altitude) at longitude  $54^\circ - 62^\circ$  east (see Figure 10). The effect spanned a range of L values and energies, but each energy channel showed the effect as being centered on a different L value (see Figure 11) in the inner radiation zone. The interpretation of Figures 10-11 is that the electron distribution was largely free of pitch-angle diffusion at longitudes east of a powerful transmitter located at about  $50^\circ - 60^\circ$  east longitude. The variation of energy with L in Figure 11 is consistent with equatorial cyclotron resonance with an unducted 16-kHz signal for reasonable plasma-density models (Vampola and Kuck, 1978).

It is tempting but dangerous to infer from such evidence that the radiation intensity in the magnetosphere was much higher in the years before those powerful VLF transmitters were installed around the world. One might argue (Vampola, 1977) that the removal of a loss mechanism inherently (and in this case dangerously) elevates particle




Figure 9. Average electron flux in the global loss cone during the year 1969 at energies 139-312 keV,  $L = 1.5-4.0$  (Vampola, 1977). Also plotted are the longitudes of VLF transmitters with authorized power levels >50 kW and >100 kW.




Figure 10. Mapping of local truncation in pitch-angle distribution to 100-km loss cone at smaller longitude (Vampola and Kuck, 1978). These OVI-19 results span a fairly wide range of L values: otherwise it would be very surprising (indeed impossible) to find such diverse truncation angles trace back to nearly the same longitude as loss-cone angles.




Figure 11. Dependence of equatorial pitch angle, corresponding to local truncation of pitch-angle distribution, upon  $L$  in the energy channel on which the effect is centered (Vampola and Kuck, 1978). For comparison, the solid curve represents the 100-km loss cone attained at a longitude of  $60^\circ$  E.



lifetimes. However, we need not conclude from the evidence that the transmitters do anything more than enhance the rate of pitch-angle diffusion at certain preferred longitudes at the expense of other longitudes. It is difficult to believe that the world's transmitters put nearly enough energy into the magnetosphere to account for the pitch-angle diffusion that occurs. The magnetosphere must have amplified their signals. This requires free energy, which is available in the anisotropy of pitch-angle distributions. Free energy in a plasma usually finds a way of expressing itself. If there are no transmitted signals to amplify, the plasma will tend to produce a broad-banded hiss through instability. On the other hand, if the free energy is spent on amplification of discrete signals, then the amount available for generating hiss will be reduced. In microscopic terms, the velocity-space diffusion caused by the amplified signals at preferred longitudes will result in a plateau on the particle distribution. The distribution thus modified is stable, and so it fails to generate hiss as the particles drift (through gradient-curvature effects) to longitudes east of the transmitter.

Actually, the whole idea that man-made transmissions account for significant particle precipitation is quite controversial. Vampola and Kuck (1978) strongly favor the idea in view of studies described above. Park and Helliwell (1978) and Lurette et al. (1979) implicitly support the idea through their studies of power-line-harmonic radiation and its effects on magnetospheric chorus. However, Tsurutani et al. (1979) argue that power-line harmonics are dynamically unimportant. Imhof et al. (1978) argue that the pitch-angle diffusion process has no significant dependence on geographic longitude. Lyons and Williams (1978) argue that anthropogenic processes act only as minor perturbations on a dynamical system that is controlled mainly by natural processes. Thus, if one is not interested in (for example) the azimuthal structure or detailed energy-latitude dependence of par-

ticle precipitation. the effect of VLF transmitters may seem unimportant since the total amount of precipitation is not greatly enhanced thereby. However, if one is interested primarily in the fine-structure of particle precipitation, then the effects of VLF transmitters and power lines may seem very important indeed.

The change in equatorial pitch angle, caused by resonance between a particle and a coherent wave packet, is strictly diffusive only in the small-amplitude limit. A numerical and analytical study by Inan et al. (1978) covers the more general case in which the wave amplitude itself is large enough to trap particles in resonance, despite the inhomogeneity of the earth's magnetic field. Such trapping is necessarily of finite duration, but it leads to a nonlinear relationship between the cumulative change  $\Delta\alpha_{eq}$  in equatorial pitch angle and the amplitude of the wave. Representative results for electrons resonant with the wave as they cross the equator are shown in Figure 12, where  $\phi_0$  denotes the initial gyrophase of the particle relative to the wave.

Results of this type are relevant for the interaction<sup>of</sup> VLF transmissions and power-line harmonics with geomagnetically trapped electrons. A related topic is the interaction of such electrons with discrete signals of time-varying frequency, e.g., whistlers and chorus elements. The nonlinear response that is evident in Figure 12 for signals  $\geq 5$  mV should serve as a warning against the uncritical application of quasilinear diffusion theory in such cases, even though the frequency spectrum appears to be smooth when averaged over time.

Other investigations of nonlinear effects have included the theoretical work by Brinca (1978) on cyclotron-resonance broadening and the numerical simulations by Papadopoulos and Rowland (1978) on the formation of auroral electron energy spectra. Moreover, Cornilleau-Wehrlin and Gendrin (1979)




Figure 12. Total change in equatorial pitch angle accumulated by electrons of various initial phases  $\phi_0$  after interaction with whistler-mode wave of specified amplitude at  $L = 4$  (Inan et al., 1978). Left panel:  $\alpha_{eq}^{(0)} = 10^\circ$ ; right panel:  $\alpha_{eq}^{(0)} = 30^\circ$ ; particle energy: that required for equatorial resonance with the 5-kHz wave.

have reported the observation of transmitter-induced quiet bands in the VLF noise spectrum and have interpreted this phenomenon in terms of nonlinear wave-particle interactions in the inhomogeneous B field. Meverson and Pokhotelov (1978), and also Meerson and Sasorov (1979), have described nonlinear bounce-resonant interactions between waves and particles. The inhomogeneity of B has likewise been important in linear-instability studies. Hasegawa (1979) has considered the bounce-resonant excitation of Alfvén waves by electron beams, while Bhatia and Lakhina (1977) have studied the drift-wave instabilities associated with the radial inhomogeneity of the ring current and plasma sheet. Buloshnikov et al. (1978) have studied proton-bounce effects on the generation of long-period geomagnetic pulsations.

The effect of an inhomogeneous B field could possibly affect linear growth rates, even on time scales short compared to the bounce period, by broadening the cyclotron resonance between waves and particles or by other equilibrium-trajectory effects (Lundin and Shkivar, 1977). The presence of an electric field parallel to B could similarly alter the linear growth rate. Ponyavin and Sazhin (1978), and also Misra et al. (1979), have studied the whistler-mode instability in the presence of such an electric field. Other investigators have focused instead on describing the Vlasov-equilibrium distribution more realistically than before. Thus, Kaufmann et al. (1978) have used the observed auroral-electron distribution (rather than a bi-Maxwellian simulation of it) for calculating the properties of electrostatic instabilities. Ashour-Abdalla and Kennel (1978a,b,c) have emphasized the effects of co-existing hot and (nearly) cold plasma components on electrostatic instabilities. Kiwamoto (1979) and Curtis and Wu (1979a,b) have emphasized the importance of gyroharmonic resonances for the case of oblique propagation.

Boswell (1978) and D'Angelo (1977) have especially considered instabilities that occur in the auroral region and polar cusp (otherwise known as the polar cleft). Finally, in work similar to (but less accurate than) the above-described work of Kaye et al. (1979), Solomon and Pellat (1978) considered the effects of magnetospheric convection on the Vlasov-equilibrium distributions of hot-plasma constituents and (consequently) on the linear growth rates of electromagnetic cyclotron waves.

## KINEMATICAL EFFECTS

It is well known that energetic outer-zone electrons have a peculiar pitch-angle distribution on the night side of the magnetosphere, such that the maximum particle flux occurs at equatorial pitch angle  $\alpha_0^*$  and  $\pi - \alpha_0^*$  different from  $\pi/2$  (e.g., West, 1979). This phenomenon was traced long ago to the properties of adiabatic drift shells in an asymmetric B field. Recently, Baker et al. (1978) have successfully used the particle anisotropy as a measure of the day-night asymmetry of the magnetosphere, i.e., as an indicator of the strength of the tail current, in an effort to predict the occurrence of geomagnetic substorms. Moreover, Luhmann and Schulz (1979) have described especially simple methods for extracting quantitative parameters from the observational data in this context. Kosik (1979) has shown how to use similar methods at particle energies for which the convection electric field must be taken into account. Whipple (1978, 1979) has demonstrated the utility of a new "coordinate" system called (U, B, K), in which the adiabatic trajectories of nonrelativistic particles are straight lines under such circumstances. The trajectories turn out to be parabolic for relativistic particles, and so the (U, B, K) system constitutes a major simplification from traditional systems even in this case.

In a study of extensive data on outer-zone electrons from three of the ATS satellites, Paulikas and Blake (1979) have shown that the particle intensity in a given energy channel correlates well with the solar-wind velocity. A portion of this effect might well be attributed to adiabatic compression of the magnetosphere and a consequent energization of the particles observed. However, a cross-correlation analysis revealed a delay  $\sim 1$ -2 days between changes in the solar-wind velocity and changes in the particle fluxes. This suggests a gradual energization of particles by processes that a

solar-wind enhancement can trigger, especially when the southward component of the interplanetary B field is positive (e.g., Akasofu, 1977). Thus, it seems that adiabatic and non-adiabatic effects (i.e., kinematical and dynamical effects) can occur in combination and thereby present a serious challenge to investigators whose duty it is to provide a quantitative description of radiation-belt processes.

## EPILOGUE

The foregoing pages offer a sample of recent developments in the study of energetic particle populations and cosmic-ray entry with respect to the earth's magnetosphere. It would not be feasible to cover all areas of progress, and the selection of topics has been largely arbitrary. However, an effort has been made to treat topics that represent significant new directions in research. Equally important progress has been made by investigators working along more traditional lines. It is inevitable in a report of this type that mention of some significant contributions to the field has been omitted, but it is hoped that the topics covered are at least representative of recent progress in this area of space research.



# REFERENCES CITED

- Akasofu, S.-I., Physics of Magnetospheric Substorms, pp. 200-257. Reidel, Dordrecht-Holland, 1977.
- Amano, K., and T. Tsuda, Particle trajectories at a magnetic neutral point. J. Geomag. Geoelectr., 30, 27, 1978.
- Ashour-Abdalla, M., and C. F. Kennel, Diffuse auroral precipitation. J. Geomag. Geoelectr., 30, 239, 1978a.
- Ashour-Abdalla, M., and C. F. Kennel, Nonconvective and convective electron cyclotron harmonic instabilities. J. Geophys. Res., 83, 1531, 1978b.
- Ashour-Abdalla, M., and C. F. Kennel, Multi-harmonic electron cyclotron instabilities. Geophys. Res. Lett., 5, 711, 1978c.
- Baker, D. N., P. R. Higbie, E. W. Hones, Jr., and R. D. Belian, High-resolution energetic particle measurements at 6.6  $R_E$ . 3. Low-energy electron anisotropies and short-term substorm predictions. J. Geophys. Res., 83, 4863, 1978.
- Baker, D. N., P. R. Higbie, R. D. Belian, and E. W. Hones, Jr., Do Jovian electrons influence the terrestrial outer radiation zone?. Geophys. Res. Lett., 6, 531, 1979.
- Belian, R. D., D. N. Baker, P. R. Higbie, and E. W. Hones, Jr., High-resolution energetic particle measurements at 6.6  $R_E$ . 2. High-energy proton drift echoes. J. Geophys. Res., 83, 4857, 1978.
- Bhatia, K. G., and G. S. Lakhina, Drift instabilities associated with the particles in ring current and inner edge of plasma sheet. Planet. Space Sci., 26, 985, 1978.
- Blake, J. B., and L. M. Friesen, A technique to determine the charge state of the anomalous low-energy cosmic rays. Paper OG-203. 15th International Cosmic Ray Conference, vol. 2, pp. 341-346, Bulg. Acad. Sci., Plovdiv, 1977.
- Blake, J. B., and L. M. Friesen, On the access and loss of few MeV protons at the geosynchronous orbit. Paper MG 2-1. 16th International Cosmic Ray Conference, vol. 3, pp. 129-133, Sci. Council and Phys. Soc. of Japan, Kyoto, 1979.
- Boswell, R. W., Energetic ions and electromagnetic radiation in auroral regions. Geophys. Res. Lett., 5, 395, 1978.
- Brinca, A. L., Turbulence effects in the cyclotron resonance of monochromatic whistlers. Geophys. Res. Lett., 5, 839, 1978.

- Buloshnikov, A. M., Ye. A. Gerasimovich, M. B. Gokhberg, and Yu. G. Khabazin. Influence of bounce effects on the generation of type Pc4,5 geomagnetic pulsations. Geomagn. Aeron., 18, 339, 1978 (Engl. translation).
- Chanteur, G., R. Gendrin, and S. Perraut. Experimental study of high-energy electron drift echoes observed on board ATS 5. J. Geophys. Res., 82, 5231, 1977.
- Chanteur, G., R. Gendrin, and S. Perraut. High-energy electron drift echoes at the geostationary orbit. J. Atmos. Terr. Phys., 40, 367, 1978.
- Cornilleau-Wehrlin, N., and R. Gendrin. VLF transmitter-induced quiet bands: A quantitative interpretation. J. Geophys. Res., 84, 882, 1979.
- Cornwall, J. M., and M. Schulz. Physics of heavy ions in the magnetosphere, in Solar System Plasma Physics, edited by L. J. Lanzerotti, C. F. Kennel, and E. N. Parker, vol. 3, pp. 165-210. North-Holland Publ. Co., Amsterdam, 1979.
- Cowley, S. W. H., A note on the motion of charged particles in one-dimensional magnetic current sheets. Planet. Space Sci., 26, 539, 1978.
- Croley, D. R., Jr., P. F. Mizera, and J. F. Fennell. Signature of parallel electric field in ion and electron distributions in velocity space. J. Geophys. Res., 83, 2701, 1978.
- Curtis, S. A., and C. S. Wu. Electrostatic and electromagnetic gyroharmonic emissions due to energetic electrons in magnetospheric plasmas. J. Geophys. Res., 84, 2057, 1979a.
- Curtis, S. A., and C. S. Wu. Gyroharmonic emissions induced by energetic ions in the equatorial plasmasphere. J. Geophys. Res., 84, 2957, 1979b.
- D'Angelo, N., Plasma waves and instabilities in the polar cusp: A review. Rev. Geophys. Space Phys., 15, 299, 1977.
- Darchieva, L. A., T. A. Ivanova, E. N. Sosnovetz, and L. V. Tverskaya. Variations of the latitude of geomagnetic cut-off of solar protons during the great geomagnetic disturbances. Geomagn. Aeron., 18, 1097, 1978 (Russian original).
- Debrunner, H., E. Flückiger, H. von Mandach, and M. Arens. Determination of the ring current radii from cosmic ray neutron monitor data for the 17 December 1971 magnetic storm. Planet. Space Sci., 27, 577, 1979.
- Frank, L. A., K. L. Ackerson, R. J. DeCoster, and B. G. Burek. Three-dimensional plasma measurements within the earth's magnetosphere. Space Sci. Rev., 22, 739, 1978.

- Gall, R., and S. Bravo, Reinterpretation of flux structures of solar protons and their use as probes of an open and closed magnetosphere. preprint, Instituto de Geofísica, Universidad Nacional Autónoma de México, México, D. F., August 1979.
- Gall, R., J. Jiménez, and L. Camacho, Arrival of low-energy cosmic rays via the magnetospheric tail, J. Geophys. Res., 73, 1593, 1968.
- Geiss, J., H. Balsiger, P. Eberhardt, H.-P. Walker, L. Weber, D. T. Young, and H. Rosenbauer, Dynamics of magnetospheric ion composition as observed by the GEOS mass spectrometer, Space Sci. Rev., 22, 537, 1978.
- Ghielmetti, A. G., R. G. Johnson, R. D. Sharp, and E. G. Shelley, The latitudinal, diurnal, and altitudinal distributions of upward flowing energetic ions of ionospheric origin, Geophys. Res. Lett., 5, 59, 1978.
- Harel, M., R. A. Wolf, P. H. Reiff, and M. Smiddy, Computer modeling of events in the inner magnetosphere, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 499-512, Am. Geophys. Union, Washington, D. C., 1979.
- Hasegawa, A., Excitation of Alfvén waves by bouncing electron beams — An origin of nightside magnetic pulsations, Geophys. Res. Lett., 6, 664, 1979.
- Holzworth, R. H., and F. S. Mozer, Direct evaluation of the radial diffusion coefficient near  $L = 6$  due to electric field fluctuations, J. Geophys. Res., 84, 2559, 1979.
- Hultqvist, B., The hot ion component of the magnetospheric plasma and some relations to the electron component — Observations and physical implications, Space Sci. Rev., 23, 581, 1979.
- Imhof, W. L., J. B. Reagan, and E. E. Gaines, The energy selective precipitation of inner zone electrons, J. Geophys. Res., 83, 4245, 1978.
- Inan, U. S., T. F. Bell, and R. A. Helliwell, Non-linear pitch-angle scattering of energetic electrons by coherent VLF waves in the magnetosphere, J. Geophys. Res., 83, 3235, 1978.
- Johnson, R. G., Energetic ion composition in the earth's magnetosphere, Rev. Geophys. Space Phys., 17, 696, 1979.
- Johnson, R. G., R. D. Sharp, and E. G. Shelley, Observations of ions of ionospheric origin in the storm-time ring current, Geophys. Res. Lett., 4, 403, 1977.
- Kaufmann, R. L., P. B. Dusenbery, and B. J. Thomas, Stability of the auroral plasma: Parallel and perpendicular propagation of electrostatic waves, J. Geophys. Res., 83, 5663, 1978.

- Kaye, S. M., M. G. Kivelson, and D. J. Southwood, Evolution of ion cyclotron instability in the plasma convection system of the magnetosphere, J. Geophys. Res., 84, 6397, 1979.
- Kiwamoto, Y., Unstable electron cyclotron harmonic waves during the relaxation of a temperature anisotropy, J. Geophys. Res., 84, 462, 1979.
- Kosik, J. C., Influence of electric fields on charged particle motion and electron fluxes at synchronous altitudes, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 569-581, Am. Geophys. Union, Washington, D. C., 1979.
- Lanzerotti, L. J., D. C. Webb, and C. W. Arthur, Geomagnetic field fluctuations at synchronous orbit, 2, Radial diffusion, J. Geophys. Res., 83, 3866, 1978.
- Luette, J. P., C. G. Park, and R. A. Helliwell, The control of magnetospheric chorus by power line radiation, J. Geophys. Res., 84, 4224, 1979 (corrected title).
- Luhmann, J. G., and M. Schulz, Magnetic shell tracing: A simplified approach, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 582-591, Am. Geophys. Union, Washington, D. C., 1979.
- Lundin, B. V., and D. R. Shklyar, Interaction of electrons with low transverse velocities with VLF waves in an inhomogeneous plasma, Geomagn. Aeron., 17, 168, 1977 (Engl. translation).
- Lyons, L. R., Generation of large-scale regions of auroral currents, electric potentials, and precipitation by the divergence of the convection electric field, J. Geophys. Res., 85, 17, 1980.
- Lyons, L. R., and D. J. Williams, A comment on the effects of man-made VLF waves on the radiation belts, Geophys. Res. Lett., 5, 116, 1978.
- Lyons, L. R., and D. J. Williams, A source for the geomagnetic storm main phase ring current, J. Geophys. Res., 85, 523, 1980.
- Meerson, B. I., and P. V. Sasorov, Nonlinear bounce resonant interaction between Alfvén waves and geomagnetically trapped particles, Planet. Space Sci., 27, 503, 1979.
- Meyerson, B. I., and O. A. Pokhotelov, Self-consistent diffusion of trapped particles on bounce-drift resonance with geomagnetic pulsations, Geomagn. Aeron., 18, 79, 1978 (Engl. translation; see reference above for alternative spelling of first author's name).
- Misra, K. D., B. D. Singh, and S. P. Mishra, Effect of a parallel electric field on the whistler mode instability in the magnetosphere, J. Geophys. Res., 84, 5923, 1979.

- Mizera, P. F., and J. F. Fennell, Signatures of electric fields from high and low altitude particle distributions, Geophys. Res. Lett., 4, 311, 1977.
- Mizera, P. F., and J. F. Fennell, Satellite observations of polar, magnetotail lobe, and interplanetary electrons at low energies, Rev. Geophys. Space Phys., 16, 147, 1978.
- Olson, W. P., A scalar potential representation of the tilted magnetopause and neutral sheet magnetic fields, Paper WD-1332, McDonnell Douglas Astronautics Co., Huntington Beach, Calif., 1970.
- Olson, W. P., and K. A. Pfitzer, A quantitative model of the magnetospheric magnetic field, J. Geophys. Res., 79, 3739, 1974.
- Park, C. G., and R. A. Helliwell, Magnetospheric effects of power line radiation, Science, 200, 727, 1978.
- Papadopoulos, K., and H. L. Rowland, Collisionless effects on the spectrum of secondary auroral electrons at low altitudes, J. Geophys. Res., 83, 5768, 1978.
- Paulikas, G. A., and J. B. Blake, Effects of the solar wind on magnetospheric dynamics: Energetic electrons at the synchronous orbit, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 180-202, Am. Geophys. Union, Washington, D. C., 1979.
- Pfitzer, K. A., The effect of magnetic field models on cosmic ray cutoff calculations, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 242-252, Am. Geophys. Union, Washington, D. C., 1979.
- Ponyavin, D. I., and S. S. Sazhin, Development of cyclotron instability of whistlers in a magnetospheric plasma with the account of a self-consistent longitudinal electric field, Geomagn. Aeron., 18, 739, 1978 (Russian original).
- Prangé, R., Energetic (keV) ions of ionospheric origin in the magnetosphere, A review, Ann. Géophys., 34, 187, 1978.
- Shelley, E. G., Heavy ions in the magnetosphere, Space Sci. Rev., 23, 465, 1979.
- Smith, P. H., N. K. Bewtra, and R. A. Hoffman, Motions of charged particles in the magnetosphere under the influence of a time-varying large scale convection electric field, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 513-535, Am. Geophys.
- Solomon, J., and R. Pellat, Convection and wave-particle interactions, J. Atmos. Terr. Phys., 40, 373, 1978.
- Spjeldvik, W. N., Expected charge states of energetic ions in the magnetosphere, Space Sci. Rev., 23, 499, 1979.

- Spjeldvik, W. N., and T. A. Fritz. Theory for charge states of energetic oxygen ions in the earth's radiation belts, J. Geophys. Res., 83, 1583, 1978a.
- Spjeldvik, W. N., and T. A. Fritz. Quiet time observations of equatorially trapped megaelectronvolt radiation belt ions with nuclear charge  $Z \geq 4$ . J. Geophys. Res., 83, 4401, 1978b.
- Swift, D. W., H. C. Stenbaek-Nielsen, and T. J. Hallinan. An equipotential model for auroral arcs, J. Geophys. Res., 81, 3931, 1976.
- Torbert, R. B., and F. S. Mozer. Electrostatic shocks as the source of discrete auroral arcs, Geophys. Res. Lett., 5, 135, 1978.
- Tsurutani, B. T., S. R. Church, and R. M. Thorne. A search for geographic control of the occurrence of magnetospheric ELF emissions, J. Geophys. Res., 84, 4116, 1979.
- Vampola, A. L., VLF transmission induced slot electron precipitation, Geophys. Res. Lett., 4, 569, 1977.
- Vampola, A. L., and G. A. Kuck. Induced precipitation of inner zone electrons. 1. Observations, J. Geophys. Res., 83, 2543, 1978.
- West, H. I., Jr., The signatures of the various regions of the outer magnetosphere in the pitch angle distributions of energetic particles, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 150-179, Am. Geophys. Union, Washington, D. C., 1979.
- Whipple, E. C., Jr., (U, B, K) coordinates: A natural system for studying magnetospheric convection, J. Geophys. Res., 83, 4318, 1978.
- Whipple, E. C., Jr., A kinetic approach to magnetospheric modeling, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 462-472, Am. Geophys. Union, Washington, D. C., 1979.
- Williams, D. J., T. A. Fritz, E. Keppler, B. Wilken, and G. Wibberenz. Three-dimensional energetic ion observations and the tangential magnetopause electric field, Space Sci. Rev., 22, 805, 1978.
- Young, D. T., Ion composition measurements in magnetospheric modeling, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 340-363, Am. Geophys. Union, Washington, D. C., 1979.

## SUPPLEMENTARY BIBLIOGRAPHY

Many more papers on energetic particle populations and cosmic-ray entry have appeared during the past two years than could be cited in the above text. The author has compiled a supplementary list of relevant works from this period, a list that is meant to be representative but not necessarily exhaustive.

Each paper has been identified with one or more general subject categories as follows: (a) inner belt; (b) outer-belt protons; (c) outer-belt electrons; (d) diffusion, wave-particle interactions; (e) injected and/or drifting particles; (f) trapped ions,  $Z \geq 2$ ; (g) solar cosmic-ray entry; (h) theory; (i) artificial perturbation of trapped particles; (j) general.

In accordance with the terrestrial orientation of IAGA Subdivision III-4, attention in the main text and in the supplementary bibliography has been restricted to the earth's magnetosphere.

Page numbers in translated journals are specified for the English (E) edition if available, otherwise for the Russian (R) edition. The list includes publications in major refereed space-science journals, as well as other materials that have come to the author's attention. No effort was made to search unrefereed journals, collections of abstracts, internal documents, and similar materials. The author apologizes for any omissions but recognizes that omissions are unavoidable in a task of this scope.

## SUPPLEMENTARY BIBLIOGRAPHY

- Akasofu, S.-I., Physics of Magnetospheric Substorms, Reidel, Dordrecht-Holland, 1977. (j)
- Aksenov, V. I., and A. V. Moshkov, Report to the XIX General Assembly of URSI. 1975-1977, Nauka, Moscow, 1978. (j)
- Alekhin, Yu. K., and N. A. Ryabova, Cyclotron interaction of a packet of VLF waves with a beam of resonance electrons, Geomagn. Aeron., 17, 691(E), 1977. (h)
- Amano, K., and T. Tsuda, Particle trajectories at a magnetic neutral point, J. Geomag. Geoelectr., 30, 27, 1978. (h)
- Anderson, K. A., R. P. Lin, F. Martel, C. S. Lin, G. K. Parks, and H. Rème, Thin sheets of energetic electrons upstream from the earth's bow shock, Geophys. Res. Lett., 6, 401, 1979. (e)
- Antonova, A. E., and N. S. Nikolaeva, Streams of energetic electrons in the outer ionosphere of the earth by observations at artificial earth satellite Prognoz-3, Geomagn. Aeron., 19, 615(R), 1979. (e)
- Antonova, E. E., and B. A. Tverskoy, About the role of polar ionosphere as a source of ions in the magnetosphere, Geomagn. Aeron., 19, 178(R), 1979. (e)
- Armstrong, T. P., S. M. Krimigis, and R. P. Lepping, Magnetosheath bursts of predominantly medium nuclei observed with Imp 8 on February 16, 1974, J. Geophys. Res., 83, 5198, 1978. (e, f)
- Arnoldy, R. L., The relationship between field-aligned current, carried by supra-thermal electrons, and the auroral arc, Geophys. Res. Lett., 4, 407, 1977. (j)
- Arnoldy, R. L., and P. B. Lewis, Jr., Correlation of ground-based and topside photometric observations with auroral electron spectra measurements at rocket altitudes, J. Geophys. Res., 82, 5563, 1977. (c)
- Arthur, C. W., J. Bjordal, and T. J. Rosenberg, Pc-3 magnetic pulsations and precipitation of energetic electrons, J. Geophys. Res., 84, 4125, 1979. (c)
- Arthur, C. W., R. L. McPherron, L. J. Lanzerotti, and D. C. Webb, Geomagnetic field fluctuations at synchronous orbit, 1. Power spectra, J. Geophys. Res., 83, 5659, 1978. (d)



- Asbridge, J. R., S. J. Bame, J. T. Gosling, G. Paschmann, and N. Sckopke, Energetic plasma ions within the earth's magnetosheath, Geophys. Res. Lett., 5, 953, 1978. (e)
- Ashour-Abdalla, M., and C. F. Kennel, Diffuse auroral precipitation, J. Geomag. Geoelectr., 30, 239, 1978. (d, h)
- Ashour-Abdalla, M., and C. F. Kennel, Nonconvective and convective electron cyclotron harmonic instabilities, J. Geophys. Res., 83, 1531, 1978. (d, h)
- Ashour-Abdalla, M., and C. F. Kennel, Multi-harmonic electron cyclotron instabilities, Geophys. Res. Lett., 5, 711, 1978. (d, h)
- Ashour-Abdalla, M., and R. M. Thorne, Toward a unified view of diffuse auroral precipitation, J. Geophys. Res., 83, 4755, 1978. (d, h)
- Baker, D. J., and H. Weil, Partially coherent radiation by charged particle beams in the ionosphere, J. Geophys. Res., 82, 5267, 1977. (d)
- Baker, D. N., R. D. Belian, P. R. Higbie, and E. W. Hones, Jr., Predictions of high-energy ( $> 0.3$  MeV) substorm-related magnetospheric particles, Proceedings of the International Solar-Terrestrial Predictions Workshop, Boulder, Colo., April 1979. (j)
- Baker, D. N., R. D. Belian, P. R. Higbie, and E. W. Hones, Jr., High-energy magnetospheric protons and their dependence on geomagnetic and interplanetary conditions, J. Geophys. Res., 84, 7138, 1979. (b)
- Baker, D. N., P. R. Higbie, R. D. Belian, and E. W. Hones, Jr., Do Jovian electrons influence the terrestrial outer radiation zone?, Geophys. Res. Lett., 6, 531, 1979. (e)
- Baker, D. N., P. R. Higbie, E. W. Hones, Jr., and R. D. Belian, High-resolution energetic particle measurements at  $6.6 R_E$ . 3. Low-energy electron anisotropies and short-term substorm predictions, J. Geophys. Res., 83, 4863, 1978. (c)
- Baker, D. N., P. R. Higbie, E. W. Hones, Jr., and R. D. Belian, The use of  $> 30$  keV electron anisotropies at  $6.6 R_E$  to predict magnetospheric substorms, Proceedings of the International Solar-Terrestrial Predictions Workshop, Boulder, Colo., April 1979. (c)
- Baker, D. N., P. Stauning, E. W. Hones, Jr., P. R. Higbie, and R. D. Belian, Strong electron pitch angle diffusion observed at geostationary orbit, Geophys. Res. Lett., 6, 205, 1979. (c)
- Baker, D. N., and E. C. Stone, The relationship of energy flow at the magnetopause to geomagnetic activity, Geophys. Res. Lett., 4, 395, 1977. (j)

- Baker, D. N., and E. C. Stone, The magnetopause energetic electron layer, 1, Observations along the distant magnetotail, J. Geophys. Res., 83, 4327, 1978. (e)
- Bame, S. J., J. R. Asbridge, J. T. Gosling, M. Haibig, G. Paschmann, N. Sckopke, and H. Rosenbauer, High temporal resolution observations of electron heating at the bow shock, Space Sci. Rev., 23, 75, 1979. (j)
- Barr, R., ELF radiation from the New Zealand power system, Planet. Space Sci., 27, 537, 1979. (i)
- Basilova, R. N., N. L. Grigorov, L. F. Kalikin, G. I. Pugacheva, and I. A. Savenko, Secondary electron fluxes at the equator, according to Kosmos-428 data, Geomagn. Aeron., 17, 225(E), 1977. (b, e)
- Basilova, R. N., E. I. Kogan-Laskina, and G. I. Pugacheva, Ionization losses of electron energy at outlet from the atmosphere with the account of change of the magnetic field strength along particle trajectory, Geomagn. Aeron., 18, 1099(R), 1978. (h)
- Basilova, R. N., Ye. I. Kogan-Laskina, and G. I. Pugacheva, Computation of ionization losses of albedo particles in the upper layers of the atmosphere with allowance for the geomagnetic field, Geomagn. Aeron., 18, 8(E), 1978. (h)
- Bassolo, V. S., and N. K. Osipov, Electrodynanic processes in the region of injection of auroral particles, Geomagn. Aeron., 17, 68(E), 1977. (j)
- Bavassano-Cattaneo, M. B., and V. Formisano, Low energy electrons and protons in the magnetosphere, Planet. Space Sci., 26, 51, 1978. (j)
- Belian, R. D., D. N. Baker, P. R. Higbie, and E. W. Hones, Jr., High-resolution energetic particle measurements at 6.6  $R_E$ , 2, High-energy proton drift echoes, J. Geophys. Res., 83, 4857, 1978. (b)
- Belov, A. V., Ya. L. Blokh, R. T. Gutschina, I. I. Dorman, E. A. Eroshenko, O. I. Inozemtseva, and N. S. Kaminer, Solar cosmic rays of 22 November 1977 by the data of the neutron component, Geomagn. Aeron., 18, 1017(R), 1978. (g)
- Belov, A. V., Ya. L. Blokh, R. T. Gutshina, I. I. Dorman, E. A. Eroshenko, O. I. Inozemtzeva, and N. S. Kaminer, Variations of cosmic rays in the period 7 to 24 September 1977 by the data of the neutron component, Geomagn. Aeron., 18, 804(R), 1978. (g)
- Benson, R. F., and W. Calvert, ISIS 1 observations at the source of auroral kilometric radiation, Geophys. Res. Lett., 6, 479, 1979. (d)
- Bernstein, W., H. Leinbach, P. Kellogg, S. Monson, I. Hallinan, O. K. Garriott, A. Konradi, J. McCoy, P. Daly, B. Baker, and H. R. Anderson, Electron beam injection experiments: The beam-plasma discharge at low pressures and magnetic field strengths, Geophys. Res. Lett., 5, 127, 1978. (i)

- Bespalov, P. A., and V. Yu. Trakhtengertz, To the theory of VLF and ELF emissions of a chorus type, Geomagn. Aeron., 18, 627(R), 1978. (d, h)
- Bhatia, K. G., and G. S. Lakhina, Generation of 1-30 Hz waves near the plasmopause by resonant electron-instability, Planet. Space Sci., 26, 195, 1978. (d, h)
- Bhatia, K. G., and G. S. Lakhina, Drift instabilities associated with the particles in ring current and inner edge of plasma sheet, Planet. Space Sci., 26, 985, 1978. (d, h)
- Birmingham, T. J., and T. G. Northrop, Theory of flux anisotropies in a guiding center plasma, J. Geophys. Res., 84, 41, 1979. (h)
- Biswas, S., and N. Durgaprasad, Skylab measurements of low energy cosmic rays, Space Sci. Rev., in press, 1980. (g)
- Bloch, G. M., and B. M. Kuzhevsky, Time variations of spectral characteristics of increase of cosmic rays before flares, Geomagn. Aeron., 19, 551(R), 1979. (g)
- Blukh, G. M., B. M. Kudzhevsky, B. M. Makhmudov, and I. A. Savenko, Investigation of relations of proton streams and  $\alpha$ -particles of small energies from solar flares, Geomagn. Aeron., 19, 609(R), 1979. (g)
- Borg, H., L.-A. Holmgren, B. Hultqvist, F. Cambou, H. Rème, A. Bahnsen, and G. Kremser, Some early results of the keV plasma experiment on GEOS-1, Space Sci. Rev., 22, 511, 1978. (j)
- Borovkov, L. P., E. V. Vashenyuk, L. L. Lazutin, and O. I. Shumilov, Solar protons in the stratosphere and at the earth's surface on May 7, 1978, Geomagn. Aeron., 19, 552(R), 1979. (g)
- Boswell, R. W., Energetic ions and electromagnetic radiation in auroral regions, Geophys. Res. Lett., 5, 395, 1978. (b, d, h)
- Brekke, A., The relationship between the Harang discontinuity and the substorm injection boundary, Planet. Space Sci., 25, 1119, 1977. (j)
- Brinca, A. L., Turbulence effects in the cyclotron resonance of monochromatic whistlers, Geophys. Res. Lett., 5, 839, 1978. (d, h)
- Brown, R. R., On sudden commencement absorption events at the south pole and their relation to the latitude of the lower cleft boundary, J. Geophys. Res., 83, 2205, 1978. (g)
- Brown, R. R., R. H. Karas, J. R. Barcus, and P. Stauning, Observations of long-period pulsations in electron precipitation at 75° magnetic latitude, J. Geophys. Res., 82, 4834, 1977. (c)

- Bryant, D. A., D. S. Hall, and D. R. Lepine, Electron acceleration in an array of auroral arcs, Planet. Space Sci., 26, 511, 1978. (a)
- Buloshnikov, A. M., and E. E. Fedorov, On the stability of "amster" and "pearl" type electromagnetic waves in the magnetosphere, Ann. Geophys., 33, 265, 1977. (d,b)
- Buloshnikov, A. M., Ye. A. Gerasimovich, M. B. Gorkberg, and Yu. L. Khabazin, Influence of bounce effects on the generation of type I-4.7 geomagnetic pulsations, Geomagn. Aeron., 15, 554-561, 1975. (a)
- Burch, J. L., S. A. Fields, and R. A. Heelis, Polar cap electron acceleration regions, J. Geophys. Res., 84, 5863, 1979. (a)
- Burch, J. L., and J. D. Winningham, Relationships between particle precipitation and auroral forms, J. Geomag. Geoelectr., 30, 1978, 1978. (j)
- Candidi, M., H. W. Kroehl, G. Kosinski, and R. W. Bunnmann, DNISF-1 observations of cusplike electron precipitation regions during the September 14-21, 1977 event, Space Sci. Rev., 22, 667, 1978. (a)
- Carbary, J. F., and S. M. Krinagis, Energetic particle activity at 5-minute and 10-second time resolution in the magnetotail and its relation to auroral activity, JHU/API preprint, Laurel, Md., December 1978. (j)
- Cartwright, D. G., S. J. Monson, and P. J. Kellogg, Heating of the ambient ionosphere by artificially injected electron beams, J. Geophys. Res., 83, 16, 1978. (i)
- Cattell, C., R. Lysak, R. B. Torbert, and F. S. Mager, Observations of differences between regions of current flowing into and out of the ionosphere, Geophys. Res. Lett., 6, 621, 1979. (a)
- Chamberlain, M. T., and F. Jacka, Optical evidence for mid-latitude charged particle precipitation, J. Atmos. Terr. Phys., 41, 111, 1979. (j)
- Chan, K., M. J. Teague, N. Schofield, and J. L. Vette, Modeling of electron time variations in the radiation belts, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 121-146, Ann. Geophys. Union, Washington, D.C., 1979. (a)
- Chanteur, G., R. Gendrin, and S. Perraut, Experimental study of 100-keV energy electron drift echoes observed on board AIS 5, J. Geophys. Res., 82, 5231, 1977. (c,e)
- Chanteur, G., R. Gendrin, and S. Perraut, High-energy electron drift echoes at the geostationary orbit, J. Atmos. Terr. Phys., 40, 161, 1978. (a)

- Charaschch'yan, T. N., Eleven-year cosmic-ray cycle in the stratosphere, Geomagn. Aeron., 18, 259(E), 1978. (e)
- Chiu, Y. T., and M. Schulz, Self-consistent particle and parallel electric field distributions in the magnetospheric-ionospheric auroral region, J. Geophys. Res., 83, 629, 1978. (j)
- Cornilleau-Wehrlin, N., and R. Gendrin, VLF transmitter-induced quiet bands: A quantitative interpretation, J. Geophys. Res., 84, 882, 1979. (i)
- Cornwall, J. M., and M. Schulz, Physics of heavy ions in the magnetosphere, in Solar System Plasma Physics, edited by C. F. Kennel, L. J. Lanzerotti, and E. N. Parker, vol. 3, pp. 165-219, North-Holland Publ. Co., Amsterdam, 1979. (f)
- Cowley, S. W. H., A note on the motion of charged particles in one-dimensional magnetic current sheets, Planet. Space Sci., 26, 539, 1978. (h)
- Croley, D. R., Jr., P. E. Mizera, and J. F. Fennell, Signature of a parallel electric field in ion and electron distributions in velocity space, J. Geophys. Res., 83, 2701, 1978. (j)
- Curtis, S. A., A theory for chorus generation by energetic electrons during substorms, J. Geophys. Res., 83, 3841, 1978. (d,h)
- Curtis, S. A., and C. S. Wu, Electrostatic and electromagnetic gyroharmonic emissions due to energetic electrons in magnetospheric plasmas, J. Geophys. Res., 84, 2057, 1979. (d,h)
- Curtis, S. A., and C. S. Wu, Gyroharmonic emissions induced by energetic ions in the equatorial plasmasphere, J. Geophys. Res., 84, 2597, 1979. (d,h)
- Curtis, S. A., C. S. Wu, and D. H. Fairfield, Electromagnetic and electrostatic emissions at the cusp-magnetosphere interface during substorms, J. Geophys. Res., 84, 898, 1979. (j)
- Davis, E. W., and B. A. Whalen, Rocket-borne measurements of particles and ion convection in dayside aurora, J. Geophys. Res., 83, 2195, 1978. (j)
- D'Angelis, N., Plasma waves and instabilities in the polar cusp: A review, Rev. Geophys. Space Phys., 15, 209, 1977. (d)
- Durand, L. A., L. A. Ivanova, E. N. Sosnovets, and L. V. Iverskaya, Variations of the latitude of geomagnetic cut-off of solar protons during the great geomagnetic disturbances, Geomagn. Aeron., 15, 106, 1978. (e)

- Davidson, G. T., Relativistic electron precipitation and resonance with ion cyclotron waves, J. Atmos. Terr. Phys., 40, 1085, 1978. (d, h)
- Debrunner, H., E. Flückiger, H. von Mandach, and M. Arens, Determination of the ring current radii from cosmic ray neutron monitor data for the 17 December 1971 magnetic storm, Planet. Space Sci., 27, 577, 1979. (g)
- DeCoster, R. J., and L. A. Frank, Observations pertaining to the dynamics of the plasma sheet, J. Geophys. Res., 84, 5099, 1979. (j)
- Domingo, V., D. E. Page, and K.-P. Wenzel, Where do charged particles enter in and exit from the magnetosphere? Some HEOS-2 measurements, J. Atmos. Terr. Phys., 40, 279, 1978. (j)
- Domingo, V., and K.-P. Wenzel, Influence of the magnetosheath on solar proton penetration into the magnetosphere, Planet. Space Sci., 25, 1111, 1977. (g)
- Duggal, S. P., Relativistic solar cosmic rays, Rev. Geophys. Space Phys., 17, 1021, 1979. (g)
- Duggal, S. P., and M. A. Pomerantz, On the source of transient cosmic ray intensity variations (Reply to comment by G. D. Parker), J. Geophys. Res., 83, 2715, 1978. (g)
- Duggal, S. P., and M. A. Pomerantz, Symmetrical equator-pole anisotropy during an unusual cosmic ray storm, Geophys. Res. Lett., 5, 625, 1978. (g)
- Durney, A. C., and K. W. Ogilvie, Introduction to the ISEE mission, Space Sci. Rev., 22, 679, 1978. (j)
- Eastman, T. E., and E. W. Hones, Jr., Characteristics of the magnetospheric boundary layer and magnetopause layer as observed by Imp 6, J. Geophys. Res., 84, 2019, 1979. (j)
- Ejiri, M., Trajectory traces of charged particles in the magnetosphere, J. Geophys. Res., 83, 4798, 1978. (e, h)
- Eliasson, L., L.-Å. Holmgren, and K. Pettersmark, Pitch angle and energy distributions of auroral electrons measured by the ESRO 4 satellite, Planet. Space Sci., 27, 87, 1979. (e)
- Elshin, V. K., G. F. Krymsky, S. I. Petukhov, Yu. A. Romaschenko, and I. A. Transky, A regular mechanism of charged particle acceleration in cosmic conditions, 1, "Superadiabatic" acceleration by shock waves, Geomagn. Aeron., 19, 606(R), 1979. (d, h)

- Erickson, K. N., R. L. Swanson, R. J. Walker, and J. R. Winckler, A study of magnetosphere dynamics during auroral electrojet events by observations of energetic electron intensity changes at synchronous orbit, J. Geophys. Res., 84, 931, 1979. (c)
- Fairfield, D. H., and E. W. Hones, Jr., Imp 6 measurements in the distant polar cusp during substorms, J. Geophys. Res., 83, 4273, 1978. (j)
- Feygin, F. Z., A. L. Kalisher, V. A. Pilipenko, O. A. Pokhotelov, and K. Dobeš, On the theory of generation of auroral radiation, Planet. Space Sci., 27, 913, 1979. (h)
- Feygin, F. Z., Ju. P. Kurchashov, V. A. Troitskaya, D. S. Fligel, and K. Dobeš, Method of locating the Pc-1 source and hot plasma parameters in the generation region, Planet. Space Sci., 27, 151, 1979. (j)
- Formisano, V., Properties of energetic electrons of magnetospheric origin in the magnetosheath and in the solar wind — correlation with geomagnetic activity, Planet. Space Sci., 27, 867, 1979. (e)
- Formisano, V., V. Domingo, and K.-P. Wenzel, On the mechanism of plasma penetration and energetic electron escape across the day-side magnetopause, J. Atmos. Terr. Phys., 40, 243, 1978. (e)
- Foster, J. C., and J. R. Burrows, Electron fluxes over the polar cap. 2. Electron trapping and energization on open field lines, J. Geophys. Res., 82, 5165, 1977. (e)
- Frank, L. A., and K. L. Ackerson, Reply to comments of E. W. Hones, Jr., J. Geophys. Res., 82, 5641, 1977. (g)
- Frank, L. A., and K. L. Ackerson, Several recent findings concerning the dynamics of the earth's magnetotail, Space Sci. Rev., 23, 375, 1978. (j)
- Frank, L. A., K. L. Ackerson, R. L. DeCoster, and B. G. Burek, Three-dimensional plasma measurements within the earth's magnetosphere, Space Sci. Rev., 23, 739, 1978. (g)
- Fraser-Smith, A. C., A weekend increase in geomagnetic activity, J. Geophys. Res., 84, 2084, 1979. (i)
- Fritz, T. A., and W. N. Spjeldvik, Observations of energetic radiation belt helium ions at the geomagnetic equator during quiet conditions, J. Geophys. Res., 83, 2574, 1978. (i)
- Fritz, T. A., and W. N. Spjeldvik, Simultaneous proton and helium ion observations of energetic radiation belt protons and helium ions at the geomagnetic equator near 1 MeV, J. Geophys. Res., 84, 2071, 1979. (i)

- Gall, R., and S. Bravo, Reinterpretation of flux structures of solar protons and their use as probes of an open and closed magnetosphere, preprint, Instituto de Geofísica, Universidad Nacional Autónoma de México, México, D. F., August 1979. (g)
- Galperin, Yu. I., V. A. Gladyshev, N. V. Jorjio, R. A. Kovrazhkin, V. M. Sinitsin, F. Cambou, J. A. Sauvaud, and J. Crasnier, Adiabatic acceleration induced by convection in the plasma sheet, J. Geophys. Res., 83, 2567, 1978. (j)
- Garrett, H. B., Review of quantitative models of the 0- to 100-keV near-earth plasma, Rev. Geophys. Space Phys., 17, 397, 1979. (j)
- Garrett, H. B., Quantitative models of the 0 to 100 keV mid-magnetospheric particle environment, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 364-388, Am. Geophys. Union, Washington, D. C., 1979. (j)
- Garrett, H. B., and S. E. DeForest, An analytical simulation of the geosynchronous plasma environment, Planet. Space Sci., 27, 1101, 1979. (j)
- Geiss, J., H. Balsiger, P. Eberhardt, H. P. Walker, L. Weber, D. T. Young, and H. Rosenbauer, Dynamics of magnetospheric ion composition as observed by the GEOS mass spectrometer, Space Sci. Rev., 22, 537, 1978. (f)
- Ghielmetti, A. G., R. G. Johnson, R. D. Sharp, and E. G. Shelley, The latitudinal, diurnal, and altitudinal distributions of upward flowing energetic ions of ionospheric origin, Geophys. Res. Lett., 5, 59, 1978. (e)
- Ghielmetti, A. G., R. D. Sharp, E. G. Shelley, and R. G. Johnson, Downward flowing ions and evidence for injection of ionospheric ions into the plasma sheet, J. Geophys. Res., 84, 5781, 1979. (e)
- Glangaud, F., A. Knob, J. Lacoume, V. Troitskaya, S. Krilov, and W. Gorbine, Movement and distribution of the energetic particles which generate Pc-1, J. Atmos. Terr. Phys., 40, 379, 1978. (d, e, h)
- Golikov, Yu. V., V. V. Pustovalov, A. B. Romanov, V. P. Silin, and V. I. Tikhonchuk, Non-linear effects of VLF noise generation by electron beams in the polar magnetosphere, Planet. Space Sci., 25, 899, 1977. (d, h)
- Gosling, J. T., J. R. Asbridge, S. J. Bame, G. Paschmann, and N. Sckopke, Observations of two distinct populations of bow shock ions in the upstream solar wind, Geophys. Res. Lett., 5, 257, 1978. (e)



- Green, J. L., D. A. Gurnett, and R. A. Hoffman, A correlation between auroral kilometric radiation and inverted V electron precipitation, J. Geophys. Res., 84, 5216, 1979. (j)
- Grigoryan, O. R., S. N. Kuznetsov, L. M. Kovrigina, E. N. Sosnovetz, and V. L. Tverskaya, Experimental definition of electron lifetimes in the outer radiation belt, Geomagn. Aeron., 19, 623(R), 1979. (c)
- Gul'yel'mi, A. V., and N. A. Zolotukhina, Generation of MHD waves of increasing frequency in the earth's magnetosphere, Geomagn. Aeron., 18, 205(E), 1978. (d)
- Gurgiolo, C., C. S. Lin, B. Mauk, G. K. Parks, and C. McIlwain, Plasma injection and diamagnetism, J. Geophys. Res., 84, 2049, 1979. (e)
- Hallinan, T. J., H. C. Stenbaek-Nielsen, and J. R. Winckler, The Echo 4 electron beam experiment: Television observation of artificial auroral streaks indicating strong beam interactions in the high-latitude magnetosphere, J. Geophys. Res., 83, 3263, 1978. (i)
- Hardy, D. A., J. W. Freeman, and H. K. Hills, Double-peaked ion spectra in the lobe plasma: Evidence for massive ions?, J. Geophys. Res., 82, 5529, 1977. (f)
- Hasegawa, A., Excitation of Alfvén waves by bouncing electron beams — An origin of nightside magnetic pulsations, Geophys. Res. Lett., 6, 664, 1979. (d, h)
- Heikkila, W. J., Comment on 'Solar wind plasma injection at the dayside magnetospheric cusp' by P. H. Reiff, T. W. Hill, and J. L. Burch, J. Geophys. Res., 83, 227, 1978. (e)
- Heikkila, W. J., R. J. Pellinen, and P. O. Welling, Energization of charged particles to high energies by an induced substorm electric field within the magnetotail, J. Geomag. Geoelectr., 30, 201, 1978. (j)
- Heinrich, W., and A. Spill, Geomagnetic shielding of cosmic rays for different satellite orbits, J. Geophys. Res., 84, 4401, 1979. (e)
- Henriksen, K., Variations of proton energy and pitch angle spectra in the upper atmosphere, J. Atmos. Terr. Phys., 41, 633, 1979. (b)
- Higbie, P. R., D. N. Baker, R. D. Belian, and E. W. Hones, Jr., Correction (reproduction of previously illegible figures), J. Geophys. Res., 84, 537, 1979. (c)
- Higbie, P. R., D. N. Baker, R. D. Belian, and E. W. Hones, Jr., Evolution of substorm and quiet-time electron anisotropies ( $30 \leq E_e \leq 300$  keV), Proceedings of the International Solar-Terrestrial Predictions Workshop, Boulder, Colo., April 1979. (c)

- Higbie, P. R., D. N. Baker, E. W. Hones, Jr., and R. D. Belian, Pitch angle distributions of  $> 30$  keV electrons at geostationary altitudes, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 203-219, Am. Geophys. Union, Washington, D.C., 1979. (c)
- Higbie, P. R., R. D. Belian, and D. N. Baker, High-resolution energetic particle measurements at  $6.6 R_E$ , 1, Electron micropulsations, J. Geophys. Res., 83, 4851, 1978. (c)
- Hill, T. W., and P. H. Reiff, Evidence of magnetospheric cusp proton acceleration by magnetic merging at the dayside magnetopause, J. Geophys. Res., 82, 3623, 1977. (b)
- Hinton, B. B., Global effects of auroral particle and Joule heating in the undisturbed thermosphere, J. Geophys. Res., 83, 707, 1978. (j)
- Holmgren, G., R. Boström, M. C. Kelley, P. M. Kintner, R. Lundin, U. V. Fahlson, E. A. Bering, and W. R. Sheldon, Artificial stimulation of auroral electron acceleration by intense field aligned currents, Geophys. Res. Lett., 6, 789, 1979. (i)
- Holzworth, R. H., and F. S. Mozer, Direct evaluation of the radial diffusion coefficient near  $L = 6$  due to electric field fluctuations, J. Geophys. Res., 84, 2559, 1979. (d)
- Hones, E. W., Jr., Substorm processes in the magnetotail: Comments on 'On hot tenuous plasmas, fireballs, and boundary layers in the earth's magnetotail' by L. A. Frank, K. L. Ackerson, and R. P. Lepping, J. Geophys. Res., 82, 5633, 1977. (j)
- Hones, E. W., Jr., Transient phenomena in the magnetotail and their relation to substorms, Space Sci. Rev., 23, 393, 1979. (j)
- Hovestadt, D., G. Gloeckler, C. Y. Fan, L. A. Fisk, F. M. Ipavich, B. Klecker, J. J. O'Gallagher, and M. Scholer, Evidence for solar wind origin of energetic heavy ions in the earth's radiation belt, Geophys. Res. Lett., 5, 1055, 1978. (f)
- Hubbard, R. F., and T. J. Birmingham, Electrostatic emissions between electron gyroharmonics in the outer magnetosphere, J. Geophys. Res., 83, 4837, 1978. (h)
- Hubbard, R. F., and G. Joyce, Simulation of auroral double layers, J. Geophys. Res., 84, 4297, 1979. (h)
- Hughes, W. J., R. L. McPherron, J. N. Barfield, and B. H. Mauk, A compressional Pc4 pulsation observed by three satellites in geostationary orbit near local midnight, Planet. Space Sci., 27, 821, 1979. (j)

- ultqvist, B., The hot ion component of the magnetospheric plasma and some relations to the electron component — Observations and physical implications, Space Sci. Rev., 23, 581, 1979. (j)
- Imhof, W. L., E. E. Gaines, and J. B. Reagan, The time persistence of certain features of electron precipitation in the slot region, J. Geophys. Res., 82, 5024, 1977. (a,c)
- Imhof, W. L., G. H. Nakano, and J. B. Reagan, Satellite observations of impulsive bremsstrahlung x-ray events associated with substorms, J. Geophys. Res., 83, 4237, 1978. (j)
- Imhof, W. L., J. B. Reagan, and E. E. Gaines, Fine-scale spatial structure in the pitch angle distributions of energetic particles near the mid-night trapping boundary, J. Geophys. Res., 82, 5215, 1977. (b,c)
- Imhof, W. L., J. B. Reagan, and E. E. Gaines, High resolution study of the spatial structure in the pitch angle distributions of energetic particles near the midnight trapping boundary, J. Geomag. Geoelectr., 30, 467, 1978. (b,c)
- Imhof, W. L., J. B. Reagan, and E. E. Gaines, The energy selective precipitation of inner zone electrons, J. Geophys. Res., 83, 4245, 1978. (a,d)
- Inan, U. S., T. F. Bell, and R. A. Helliwell, Nonlinear pitch angle scattering of energetic electrons by coherent VLF waves in the magnetosphere, J. Geophys. Res., 83, 3235, 1978. (h,i)
- Intriligator, D. S., H. R. Collard, J. D. Mihalov, O. L. Vaisberg, and J. H. Wolfe, Evidence for earth magnetospheric tail associated phenomena at 3100 R<sub>E</sub> Geophys. Res. Lett., 6, 585, 1979. (j)
- Ipavich, F. M., G. Gloeckler, C. Y. Fan, L. A. Fisk, D. Hovestadt, B. Klecker, J. J. O'Gallagher, and M. Scholer, Initial observations of low energy charged particles near the earth's bow shock on ISEE-1, Space Sci. Rev., 23, 93, 1979. (j)
- Izhovkina, N. I., Parameters of the albedo electron flux for electron pulses artificially injected into the ionosphere, Geomagn. Aeron., 18, 246(E), 1978. (i)
- Izhovkina, N. I., Spreading of the distribution function of an electron beam in the ionosphere as a function of the distance to the injection point, Geomagn. Aeron., 18, 353(E), 1978. (e)
- Izhovkina, N. I., Brightness of artificial aurora depending on pitch angle of injection of an electron impulse, Geomagn. Aeron., 19, 176(R), 1979. (i)
- Izhovkina, N. I., Secondary effects in interaction of an artificially injected electron beam with neutrals of the atmosphere in near-rocket space, Geomagn. Aeron., 19, 371(R), 1979. (i)

- Izhovkina, I. N., and E. V. Mishin, About a possibility of lightning to a plasma-bunch discharge when auroral electrons penetrate into the ionosphere, Geomagn. Aeron., 19, 585(R), 1979. (j)
- Janni, J. F., and G. E. Radke, The radiation environment and its effects on spacecraft, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 634-643, Am. Geophys. Union, Washington, D.C., 1979. (j)
- Jayanthi, U. B., Composition and energy spectrum of low energy carbon to nickel ions at Skylab altitudes: Sampling of the outer radiation belt? Ann. Géophys., 35, 27, 1979. (f)
- Jentsch, V., and G. Wibberenz, An analytic study of the energy and pitch angle distribution of inner-zone protons, preprint, Institut für Reine und Angewandte Kernphysik, Universität Kiel, June 1979. (a)
- Johnson, R. G., Energetic ion composition in the earth's magnetosphere, Rev. Geophys. Space Phys., 17, 696, 1979. (f)
- Johnson, R. G., R. D. Sharp, and E. G. Shelley, Observations of ions of ionospheric origin in the storm-time ring current, Geophys. Res. Lett., 4, 403, 1977. (b,e)
- Johnstone, A. D., Precipitation of charged particles by a parallel electric field, Planet. Space Sci., 26, 581, 1978. (j)
- Kamide, Y., and J. D. Winningham, A statistical study of the 'instantaneous' nightside auroral oval: The equatorward boundary of electron precipitation as observed by the Isis 1 and 2 satellites, J. Geophys. Res., 82, 5573, 1977. (c)
- Kan, J. R., L. C. Lee, and S.-I. Akasofu, Two-dimensional potential double layers and discrete auroras, J. Geophys. Res., 84, 4305, 1979. (j)
- Kasimov, U., and V. P. Shabansky, Longitudinal transformation of the density of captured particles in remote regions of the magnetosphere, Geomagn. Aeron., 19, 490(R), 1979. (e)
- Kaufmann, R. L., P. B. Dusenbery, B. J. Thomas, and R. L. Arnoldy, Auroral electron distribution function, J. Geophys. Res., 83, 586, 1978. (c)
- Kaufmann, R. L., P. B. Dusenbery, and B. J. Thomas, Stability of the auroral plasma: Parallel and perpendicular propagation of electrostatic waves, J. Geophys. Res., 83, 5663, 1978. (d,h)
- Kaye, S. M., and M. G. Kivelson, Time dependent convection electric fields and plasma injection, J. Geophys. Res., 84, 4183, 1979. (e,h)

- Kaye, S. M., C. S. Lin, G. K. Parks, and J. R. Winckler, Adiabatic modulation of equatorial pitch angle anisotropy, J. Geophys. Res., 83, 2675, 1978. (c)
- Kellogg, P. J., S. J. Monson, and B. A. Whalen, Rocket observation of high frequency waves over a strong aurora, Geophys. Res. Lett., 5, 47, 1978. (j)
- Khabazin, Yu. G., O. A. Pokhotelov, and A. V. Forstov, The role of proton injection processes in the nonlinear evolution of "pearls," Planet. Space Sci., 27, 165, 1979. (d,e,h)
- Khor, H. P., W. K. Kwok, and A. J. Owens, The annual and semiannual variations of the cosmic radiation, J. Geophys. Res., 84, 5272, 1979. (e)
- Kintner, P. M., K. L. Ackerson, D. A. Gurnett, and L. A. Frank, Correlated electric field and low-energy electron measurements in the low-altitude polar cusp, J. Geophys. Res., 83, 163, 1978. (c)
- Kintner, P. M., M. C. Kelley, and F. S. Mozer, Electrostatic hydrogen cyclotron waves near one earth radius altitude in the polar magnetosphere, Geophys. Res. Lett., 5, 139, 1978. (d)
- Kirsch, E., M. Scholer, and L. Rossmberg, Substorm related electron and proton bursts over the polar caps, Planet. Space Sci., 25, 431, 1977. (b,c)
- Kisabeth, J. L., and G. Rostoker, Relationship of noise in the frequency range  $100 < f < 500$  kHz to auroral arcs and field-aligned current and implications regarding acceleration of auroral electrons, J. Geophys. Res., 84, 853, 1979. (j)
- Kiwamoto, Y., Unstable electron cyclotron harmonic waves during the relaxation of a temperature anisotropy, J. Geophys. Res., 84, 462, 1979. (h)
- Klumppar, D. M., Transversely accelerated ions: An ionospheric source of hot magnetospheric ions, J. Geophys. Res., 84, 4229, 1979. (b)
- Knoll, G., K. Holzer, and G. Schmidtke, Spatial shift of the South Atlantic anomaly, J. Geophys. Res., 82, 5281, 1977. (j)
- Knott, K., The GEOS-1 mission, Space Sci. Rev., 22, 321, 1978. (c)
- Kohno, T., and M. Wada, Unusual arrival times of scatter-free particles in solar proton event of May 7, 1978, Geophys. Res. Lett., 6, 421, 1979. (g)

- Kolchin, A. A., V. V. Lebedev, V. F. Levchenko, A. I. Repin, G. P. Serebtzov, and V. L. Shubin, Enrichment by heavy nuclei of small energies in the solar cosmic rays on October 12-15, 1977, by the data of the experiment Ayaks-M at the automatic station Prognoz-6, Geomagn. Aeron., 19, 351(R), 1979. (g)
- Koons, H. C., M. H. Dazey, and B. C. Edgar, Satellite observation of discrete VLF line radiation within transmitter-induced amplification bands, J. Geophys. Res., 83, 3887, 1978. (i)
- Koons, H. C., and M. B. Pongratz, Ion cyclotron waves generated by an ionospheric barium injection, J. Geophys. Res., 84, 533, 1979. (ii)
- Korth, A., G. Kremser, and B. Wilken, Observations of substorm-associated particle-flux variations at  $6 \leq L \leq 8$  with GEOS-1, Space Sci. Rev., 22, 501, 1978. (b,c)
- Kosik, J. C., The uses of past and present magnetospheric field models for mapping fluxes and calculating conjugate points, Space Sci. Rev., 22, 481, 1978. (e,h)
- Kosik, J. C., Influence of electric fields on charged particle motion and electron fluxes at synchronous altitudes, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 560-581, Am. Geophys. Union, Washington, D. C., 1979. (e,h)
- Kovner, M. S., V. A. Kuznetsova, and Ya. I. Likhter, On the mid-latitude modulation of fluxes of energetic electrons and VLF emission Geomagn. Aeron., 17, 575(E), 1977. (c,d)
- Kovrygina, L. M., E. N. Sosnovets, and L. V. Iverskaya, Formation and decay of the belt of energetic electrons injected during the storm of July 4-6, 1974, Geomagn. Aeron., 17, 351(E), 1977. (c,e)
- Kovrigina, L. M., and L. V. Iverskaya, About injection of relativistic electrons into radiation belts during magnetic storms, Geomagn. Aeron., 18, 749(R), 1978. (c,e)
- Kremser, G., H. Specht, E. Kirsch, K. H. Saeger, and W. Riedler, Energetic solar-proton precipitation in the auroral zone associated with storm sudden commencements, Planet. Space Sci., 25, 823, 1977. (g)
- Krimigis, S. M., and E. I. Sarris, Energetic particle bursts in the earth's magnetotail, JHU/API preprint, Laurel, Md., December 1978. (ii)
- Krimigis, S. M., D. Venkatesan, L. C. Barichello, and E. I. Sarris, Simultaneous measurements of energetic protons and electrons in the distant magnetosheath, magnetotail, and upstream in the solar wind, Geophys. Res. Lett., 5, 301, 1978. (b,c,g)

- Krimsky, P. F., V. G. Moiseev, N. G. Skryabin, and V. D. Sokolov, Dependence of profile of temporal distribution of a stream of outpouring particles from their pitch-angle distribution, Geomagn. Aeron., 19, 691(R), 1979. (j)
- Kropotkin, A. P., Evolution of energetic-particle streams upon injection into the magnetosphere, Geomagn. Aeron., 17, 175(E), 1977. (g)
- Kropotkin, A. P., About the nature of longitudinal electric fields on auroral force lines, Geomagn. Aeron., 19, 686(R), 1979. (j)
- Kudela, K., B. Dobrovolska, A. V. Zakharov, and V. A. Kuznetsova, 440 keV proton precipitation at middle latitudes during the recovery phase of magnetic storms, Planet. Space Sci., 25, 1186, 1977. (b)
- Kurth, W. S., M. Ashour-Abdalla, L. A. Frank, C. F. Kennel, D. A. Gurnett, D. D. Sentman, and B. G. Burek, A Comparison of intense electrostatic waves near  $f_{CHR}$  with linear instability theory, Geophys. Res. Lett., 6, 487, 1979. (d, h)
- Kuznetsov, S. N., L. L. Lazutin, and V. Ye. Tsirs, Dynamics of trapped electrons in the auroral zone during a substorm, Geomagn. Aeron., 17, 180(E), 1977. (c)
- Kuznetsov, S. N., G. B. Lopatina, and V. G. Stolpovskiy, Outer boundary of the radiation belts, according to Kosmos-426 data, Geomagn. Aeron., 17, 274(E), 1977. (j)
- Kyker, G. C., Jr., and A. R. Liboff, Absolute cosmic ray ionization measurements in a 900-liter chamber, J. Geophys. Res., 83, 5539, 1978. (g)
- Lanzerotti, L. J., D. C. Webb, and C. W. Arthur, Geomagnetic field fluctuations at synchronous orbit, 2. Radial diffusion, J. Geophys. Res., 83, 3866, 1978. (d)
- Lassen, K., The quiet-time pattern of auroral arcs as a consequence of magnetospheric convection, Geophys. Res. Lett., 6, 777, 1979. (g)
- Lazarev, V. I., L. V. Iverskaya, M. V. Tel'tsov, and O. V. Khorosheva, Asymmetric injection of ring-current protons during the storm of July 6, 1974, Geomagn. Aeron., 17, 110(E), 1977. (b, e)
- Lazutin, L. L., and V. E. Tsirs, Fresh energetic electron precipitation at the noon sector of the auroral magnetosphere during the recovery phase of substorms, J. Geomag. Geoelectr., 30, 203, 1978. (c)
- Leinbach, H., and Williams, D. L., Evidence for very weak pitch angle diffusion of outer zone electrons, J. Geophys. Res., 82, 5091, 1977. (c)

- Lemaire, J., Impulsive penetration of filamentary plasma elements into the magnetospheres of the earth and Jupiter, Planet. Space Sci., 25, 887, 1977. (e)
- Lemaire, J., M. J. Rycroft, and M. Roth, Control of impulsive penetration of solar wind irregularities into the magnetosphere by the interplanetary magnetic field direction, Planet. Space Sci., 27, 47, 1979. (e)
- Lennartsson, W., E. G. Shelley, R. D. Sharp, R. G. Johnson, and H. Balsiger, Some initial ISEE-1 results on the ring current composition and dynamics during the magnetic storm of December 11, 1977, Geophys. Res. Lett., 6, 483, 1979. (f)
- Lin, C. S., and R. A. Hoffman, Characteristics of the inverted-V event, J. Geophys. Res., 84, 1514, 1979. (j)
- Lin, C. S., and R. A. Hoffman, Fluctuations of inverted V electron fluxes, NASA/GSFC preprint, Greenbelt, Md., July 1979. (c)
- Lin, C. S., B. Mauk, G. K. Parks, S. DeForest, and C. E. McIlwain, Temperature characteristics of electron beams and ambient particles, J. Geophys. Res., 84, 2651, 1979. (j)
- Lockwood, J. A., L. Hsieh, L. Friling, C. Chen, and D. Swartz, Atmospheric neutron and gamma ray fluxes and energy spectra, J. Geophys. Res., 84, 1402, 1979. (j)
- Lohrey, B., and A. B. Kaiser, Whistler-induced anomalies in VLF propagation, J. Geophys. Res., 84, 5121, 1979. (d)
- Lurette, J. P., C. G. Park, and R. A. Helliwell, The control of the magnetosphere by power line radiation (corrected title: The control of magnetospheric chorus by power line radiation), J. Geophys. Res., 84, 2657, 1979. (i)
- Luhmann, J. G., Auroral pulsations from atmospheric waves, J. Geophys. Res., 84, 4224, 1979. (j)
- Luhmann, J. G., and J. B. Blake, Rocket altitude atmospheric x-rays from magnetospheric electrons at low and middle latitudes, J. Atmos. Terr. Phys., 40, 465, 1978. (a,c)
- Luhmann, J. G., and J. B. Blake, K-shell x-ray fluxes at satellite altitude from proton precipitation, J. Atmos. Terr. Phys., 40, 471, 1978. (a,b)
- Luhmann, J. G., H. R. Rugge, J. B. Blake, and L. A. Christopher, Low latitude atmospheric x-rays observed by HEAO-1, Geophys. Res. Lett., 6, 25, 1979. (a)
- Luhmann, J. G., and M. Schulz, Magnetic shell tracing: A simplified approach, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 582-591, Am. Geophys. Union, Washington, D.C., 1979. (e,h)



- Lui, A. T. Y., L. A. Frank, K. L. Ackerson, C.-I. Meng, and S.-I. Akasofu, Systematic study of plasma flow during plasma sheet thinnings, J. Geophys. Res., 82, 4815, 1977. (e)
- Lui, A. T. Y., and C.-I. Meng, Relevance of southward magnetic fields in the neutral sheet to anisotropic distribution of energetic electrons and substorm activity, J. Geophys. Res., 84, 5817, 1979. (c)
- Lukkari, L., J. Kangas, and H. Ranta, Correlated electron precipitation and magnetic IPDP events near the plasmapause, J. Geophys. Res., 82, 4750, 1977. (c)
- Lundblad, J. Å., and F. Søråas, Proton observations supporting the ion cyclotron wave heating theory of SAR arc formation, Planet. Space Sci., 26, 245, 1978. (b,d)
- Lundblad, J. Å., F. Søråas, and K. Aarsnes, Substorm morphology of >100 keV protons, Planet. Space Sci., 27, 841, 1979. (b)
- Lundin, B. V., and D. R. Shklyar, Interaction of electrons with low transverse velocities with VLF waves in an inhomogeneous plasma, Geomagn. Aeron., 17, 168(E), 1977. (d,h)
- Lyons, L. R., Magnetic field distortions at  $L = 3-4$  inferred from simultaneous low and high altitude observations of ring current ions, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 64-76, Am. Geophys. Union, Washington, D.C., 1979. (b)
- Lyons, L. R., Plasma processes in the earth's radiation belts, in Solar System Plasma Physics, edited by C. F. Kennel, L. J. Lanzerotti, and E. N. Parker, vol. 3, pp. 137-163, North-Holland Publ. Co., Amsterdam, 1979. (j)
- Lyons, L. R., Generation of large-scale regions of auroral currents, electric potentials, and precipitation by the divergence of the convection electric field, NOAA/SEL preprint, Boulder, Colo., July 1979. (j)
- Lyons, L. R., D. S. Evans, and R. Lundin, An observed relation between magnetic field aligned electric fields and downward electron energy fluxes in the vicinity of auroral forms, J. Geophys. Res., 84, 457, 1979. (c)
- Lyons, L. R., and A. D. Richmond, Low-latitude E region ionization by energetic ring current particles, J. Geophys. Res., 83, 2201, 1978. (b)
- Lyons, L. R., and D. J. Williams, A comment on the effects of man-made VLF waves on the radiation belts, Geophys. Res. Lett., 5, 116, 1978. (i)

- Lyons, L. R., and D. J. Williams, A source for the geomagnetic storm main phase ring current, NOAA/SEL preprint, Boulder, Colo., February 1979. (b,f)
- Lysak, R. L., and M. K. Hudson, Coherent anomalous resistivity in the region of electrostatic shocks, Geophys. Res. Lett., 6, 661, 1979. (h)
- MacLennan, C. G., L. J. Lanzerotti, A. Hasegawa, E. A. Bering, III, J. R. Benbrook, W. R. Sheldon, T. J. Rosenberg, and D. L. Matthews, On the relationship of  $\sim 3$  mHz (Pc-5) electric, magnetic, and particle variations, Geophys. Res. Lett., 5, 403, 1978. (j)
- Maeda, K., N. K. Bewtra, and P. H. Smith, Ring current electron trajectories associated with VLF emissions, J. Geophys. Res., 83, 4339, 1978. (c,d,e)
- Maggs, J. E., Electrostatic noise generated by the auroral electron beam, J. Geophys. Res., 83, 3173, 1978. (d,h)
- Managadze, G. G., and V. V. Pletnev, Energy distribution of the electrons reflected from the conjugate region in the ARAKS experiment, Soviet J. Plasma Phys., 5, 89(E), 1979. (i)
- Matthews, T., S. P. Agrawal, and D. Venkatesan, A comparison of ground level cosmic ray enhancements of November 22, 1977, and February 25, 1969, Geophys. Res. Lett., 5, 1063, 1978. (g)
- McDiarmid, I. B., J. R. Burrows, and M. D. Wilson, Comparison of magnetic field perturbations at high latitudes with charged particle and IMF measurements, J. Geophys. Res., 83, 681, 1978. (j)
- McDiarmid, I. B., J. R. Burrows, and M. D. Wilson, Large-scale magnetic field perturbations and particle measurements at 1400 km on the dayside, J. Geophys. Res., 84, 1431, 1979. (j)
- McDiarmid, D. R., and A. B. McNamara, Radio aurora in the dayside auroral oval, 1. Spatial relationship with field-aligned currents and energetic particles, J. Geophys. Res., 83, 3226, 1978. (j)
- McEwen, D. J., Electron precipitation observations from a rocket flight through the dayside auroral oval, Planet. Space Sci., 25, 1161, 1977. (c)
- McEwen, D. J., and D. A. Bryant, Optical-particle characteristics of pulsating aurora, J. Atmos. Terr. Phys., 40, 871, 1978. (j)
- McEwen, D. J., and P. Venkatarangan, Electron flux and auroral intensity measurements in situ, Geophys. Res. Lett., 5, 1051, 1978. (c)
- Meerson, B. I., and P. V. Sasorov, Nonlinear bounce resonant interaction between Alfvén waves and geomagnetically trapped particles, Planet. Space Sci., 27, 503, 1979. (d,h)

- Meng, C.-I., Electron precipitations and polar auroras, Space Sci. Rev., 22, 223, 1978. (c)
- Meng, C.-I., Conjugate low energy electron observations made by ATS-6 and DMSP-32 satellites, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 96-109, Am.Geophys. Union, Washington, D.C., 1979. (c)
- Meng, C.-I., S.-I. Akasofu, and K. A. Anderson, Dawn-dusk gradient of the precipitation of low-energy electrons over the polar cap and its relation to the interplanetary magnetic field, J. Geophys. Res., 82, 5271, 1977. (e)
- Meng, C.-I., B. Mauk, and C. E. McIlwain, Electron precipitation of evening diffuse aurora and its conjugate electron fluxes near the magnetospheric equator, J. Geophys. Res., 84, 2545, 1979. (c)
- Meng, C.-I., A. L. Snyder, Jr., and H. W. Kroehl, Observations of westward traveling surges and electron precipitation, J. Geophys. Res., 83, 575, 1978. (c)
- Mewaldt, R. A., E. C. Stone, and R. E. Vogt, Characteristics of the spectra of protons and alpha particles in recurrent events at 1 AU, Geophys. Res. Lett., 6, 589, 1979. (g)
- Meyerson, B. I., and O. A. Pokhotelov, Self-consistent diffusion of trapped particles on bounce-drift resonance with geomagnetic pulsations, Geomagn. Aeron., 18, 79(E), 1978. (d,h)
- Mikerina, N. V., A. V. Belov, and K. G. Ivanov, A geomagnetic storm and variations of galactic cosmic rays when passing a burst stream with a complicated hydromagnetic structure, Geomagn. Aeron., 19, 439(R), 1979. (e)
- Mikirova, N. A., and N. K. Pereyaslova, Peculiarities of proton penetration in the auroral regions, Geomagn. Aeron., 18, 728(R), 1978. (b)
- Mineev, Yu. V., I. A. Savenko, V. G. Saveliev, and E. S. Spir'kova, Differential energy spectra and electron bursts near the magnetosphere, Geomagn. Aeron., 19, 154(R), 1979. (e)
- Mineyev, Yu. V., I. A. Savenko, V. G. Savel'yev, and Ye. S. Spir'kova, Study of electrons with energies of 0.3-3 MeV on the Prognoz-4 satellite, Geomagn. Aeron., 18, 138(E), 1978. (c)
- Misra, K. D., B. D. Singh, and S. P. Mishra, Effect of a parallel electric field on the whistler mode instability in the magnetosphere, J. Geophys. Res., 84, 5923, 1979. (d,h)
- Mizera, P. T., J. G. Luhmann, W. A. Kolasinski, and L. B. Blake, Correlated observations of auroral arcs, electrons, and X rays from a DMSP satellite, J. Geophys. Res., 83, 5573, 1978. (g)

- Mizera, P. F., and J. F. Fennell, Satellite observations of polar, magnetotail lobe, and interplanetary electrons at low energies, Rev. Geophys. Space Phys., 16, 147, 1978. (e)
- Mizera, P. F., and J. F. Fennell, Signatures of electric fields from high and low altitude particle distributions, Geophys. Res. Lett., 4, 311, 1977. (b,c)
- Mori, S., and K. Nagashima, Inference of sector polarity of the interplanetary magnetic field from the cosmic ray north-south asymmetry, Planet. Space Sci., 27, 39, 1979. (g)
- Murphree, J. S., and H. R. Anderson, Frequency analysis of 4- to 6-keV electrons associated with an auroral arc, J. Geophys. Res., 83, 730, 1978. (c)
- Murphree, J. S., and C. D. Anger, Instantaneous auroral particle energy deposition as determined by optical emission, Geophys. Res. Lett., 5, 551, 1978. (j)
- Nakano, G. H., W. L. Imhof, and J. B. Reagan, Coherent microburst electron precipitation measured from a satellite, J. Geomag. Geoelectr., 30, 347, 1978. (c)
- Nishida, A., Radial diffusion equation for collision-affected plasma, Geophys. Res. Lett., 5, 63, 1978. (d,h)
- Novikov, A. M., Yu. G. Shafer, and V. D. Sokolov, Relation between auroral x-rays and the VLF emission of the upper atmosphere, Geomagn. Aeron., 17, 199(E), 1977. (c,d)
- O'Brien, K., Secular variations in the production of cosmogenic isotopes in the earth's atmosphere, J. Geophys. Res., 84, 423, 1979. (j)
- O'Brien, K., H. A. Sandmeier, G. E. Hansen, and J. F. Campbell, Cosmic ray induced neutron background sources for geometries of air over water, ground, iron, and aluminum, J. Geophys. Res., 83, 114, 1978. (h)
- Ogilvie, K. W., and J. D. Scudder, First results from the six-axis electron spectrometer on ISEE-1, Space Sci. Rev., 23, 123, 1979. (c)
- Ogilvie, K. W., T. von Rosenvinge, and A. C. Durney, International sun-earth explorer: A three-spacecraft program, Science, 198, 131, 1977. (j)
- Okhlopkov, V. P., L. S. Okhlopkova, and T. N. Charakhchyan, Two-year variations of cosmic rays, Geomagn. Aeron., 19, 431(R), 1978. (e)
- O'Neil, R. R., F. Bien, D. Burt, J. A. Sandock, and A. T. Starr, et al., Summarized results of the artificial auroral experiment, Geophys. Res. Lett., 5, 551, 1978. (j)

- O'Neil, R. R., O. Shepherd, W. P. Reidy, J. W. Carpenter, I. M. Davis, D. Newell, J. C. Ulwick, and A. T. Stair, Jr., Exceeds 2 test, an artificial auroral experiment: Ground-based optical measurements. J. Geophys. Res., 83, 3281, 1978. (i)
- Owens, A. J., The effects of fluctuations and noise on the neutron monitor diurnal anisotropy, 2, Non-field-aligned diffusion. J. Geophys. Res., 82, 5551, 1977. (j)
- Owens, A. J., The effects of fluctuation and noise on the neutron monitor diurnal anisotropy, 3, Total vector anisotropy. J. Geophys. Res., 84, 125, 1979. (j)
- Palmer, I. D., and G. S. Craig, Distortion of interplanetary pitch distributions by earth's bow shock, J. Geophys. Res., 83, 5633, 1978. (e)
- Palmer, I. D., and P. R. Higbie, Magnetosheath distortion of pitch angle distributions of solar protons, J. Geophys. Res., 83, 30, 1978. (g)
- Palmer, I. D., and E. W. Hones, Jr., Characteristics of energetic electrons in the vicinity of the magnetospheric boundary layer at Vela orbit. J. Geophys. Res., 83, 2584, 1978. (e)
- Palmer, I. D., J. C. Jackson, and E. W. Hones, Jr., The solar proton event of April 16, 1970, 3, Evolution of pitch angle distributions as  $\leq 1$ -MeV protons propagate into the high-latitude magnetotail. J. Geophys. Res., 84, 109, 1979. (g)
- Palmer, I. D., J. C. Jackson, and E. W. Hones, Jr., Entry of solar protons to the plasma sheet and lobe of the magnetotail at  $r = 16 R_E$  in the event of April 22, 1971. J. Geophys. Res., 84, 2630, 1979. (g)
- Papadopoulos, K., and H. L. Rowland, Collisionless effects on the spectrum of secondary auroral electrons at low altitudes. J. Geophys. Res., 83, 5768, 1978. (d,h)
- Park, C. G., and D. C. D. Chang, Transmitter simulation of power line radiation effects in the magnetosphere. Geophys. Res. Lett., 5, 861, 1978. (i)
- Park, C. G., and R. A. Helliwell, Whistler precursors: A possible catalytic role of power line radiation. J. Geophys. Res., 82, 3634, 1977. (d,i)
- Park, C. G., and T. R. Miller, Sunday decreases in magnetospheric VLF wave activity. J. Geophys. Res., 84, 943, 1979. (i)
- Parker, G. D., Comment on 'The origin of transient cosmic ray intensity variations' by S. P. Duggal and M. A. Pomerantz. J. Geophys. Res., 83, 2711, 1978. (g)
- Parks, G. K., Microburst precipitation phenomena. J. Geomag. Geoelectr., 30, 327, 1978. (c)

- Parks, G. K., C. Gurgiolo, C. S. Lin, K. A. Anderson, K. P. Lin, F. Martel, and H. Rème, Dual spacecraft observations of energetic particles in the vicinity of the magnetopause, bow shock, and the interplanetary medium, Space Sci. Rev., 22, 765, 1978. (e)
- Parks, G. K., C. Gurgiolo, and R. West, Relativistic electron precipitation, Geophys. Res. Lett., 6, 393, 1979. (c)
- Parks, G. K., C. S. Lin, B. Mauk, S. DeForest, and C. E. McIlwain, Characteristics of magnetospheric particle injection deduced from events on August 18, 1974, J. Geophys. Res., 82, 5208, 1977. (e)
- Paschmann, G., G. Haerendel, N. Sckopke, and H. Rosenbauer, The entry layer, J. Atmos. Terr. Phys., 40, 257, 1978. (e)
- Paschmann, G., N. Sckopke, S. J. Bame, J. R. Asbridge, J. T. Gosling, C. T. Russell, and E. W. Greenstadt, Association of low-frequency waves with suprathermal ions in the upstream solar wind, Geophys. Res. Lett., 6, 209, 1979. (d)
- Paschmann, G., N. Sckopke, G. Haerendel, J. Papamastorakis, S. J. Bame, J. R. Asbridge, J. T. Gosling, E. W. Hones, Jr., and E. R. Tech, ISEE plasma observations near the subsolar magnetopause, Space Sci. Rev., 22, 717, 1978. (j)
- Paulikas, G. A., The space radiation environment: Effects on space systems, in Solar System Plasma Physics, edited by C. F. Kennel, L. J. Lanzerotti, and E. N. Parker, vol. 3, pp. 330-336, North-Holland Publ. Co., Amsterdam, 1979. (j)
- Paulikas, G. A., and J. B. Blake, Effects of the solar wind on magnetospheric dynamics: Energetic electrons at the synchronous orbit, Aerospace Report No. ATR-79(7642)-1, El Segundo, Calif., November 1978. (c)
- Paulikas, G. A., and J. B. Blake, Effects of the solar wind on magnetospheric dynamics: Energetic electrons at the synchronous orbit, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 180-202, Am. Geophys. Union, Washington, D.C., 1979. (c)
- Pellinen, R. J., and W. J. Heikkila, Energization of charged particles to high energies by an induced substorm electric field within the magnetotail, J. Geophys. Res., 83, 1544, 1978. (h)
- Penman, J. M., J. K. Hargreaves, and C. E. McIlwain, The relation between 10 to 80 keV electron precipitation observed at geosynchronous orbit and auroral radio absorption observed with riometers, Planet. Space Sci., 27, 445, 1979. (c)
- Peteiski, E. F., U. Fahleson, and S. D. Shawhan, Models for quasi-periodic electric fields and associated electron precipitation in the auroral zone, J. Geophys. Res., 83, 2489, 1978. (c)

- Peterson, W. K., E. G. Shelley, R. D. Sharp, R. G. Johnson, J. Geiss, and H. Rosenbauer,  $H^+$  and  $He^{++}$  in the dawnside magnetosheath, Geophys. Res. Lett., 6, 667, 1979. (j)
- Pfitzer, K. A., The effect of magnetic field models on cosmic ray cutoff calculations, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 242-252, Am. Geophys. Union, Washington, D. C., 1979. (g,h)
- Pilipenko, V. A., and O. A. Pokhotelov, Drift-mirror instability in a curved magnetic field, Geomagn. Aeron., 17, 112(E), 1977. (j)
- Pilipp, W. G., and G. Morfill, The formation of the plasma sheet resulting from plasma mantle dynamics, J. Geophys. Res., 83, 5670, 1978. (e)
- Pirog, O. M., and V. D. Urbanovitch, A model of interaction of electron streams with the atmosphere, Geomagn. Aeron., 18, 610(R), 1978. (h)
- Pokhotelov, O. A., and Yu. G. Khabazin, Influence of a geomagnetic field distortion by solar wind pressure on development of cyclotron instability of fast magnetosound waves in a magnetospheric plasma, Geomagn. Aeron., 18, 1077(R), 1978. (h)
- Pomerantz, M. A., and S. P. Duggal, Relativistic solar cosmic rays during the current sunspot cycle, Geophys. Res. Lett., 5, 877, 1978. (g)
- Ponyavin, D. I., and S. S. Sazhin, Development of cyclotron instability of whistlers in a magnetospheric plasma with the account of a self-consistent longitudinal electric field, Geomagn. Aeron., 18, 739(R), 1978. (d,h)
- Potemra, T. A., J. P. Doering, W. K. Peterson, C. O. Bostrom, R. A. Hoffman, and L. H. Brace, AE-C observations of low-energy particles and ionospheric temperatures in the turbulent polar cusp: Evidence for the Kelvin-Helmholtz instability, J. Geophys. Res., 83, 3877, 1978. (e)
- Potemra, T. A., W. K. Peterson, J. P. Doering, C. O. Bostrom, R. W. McEntire, and R. A. Hoffman, Low-energy particle observations in the quiet dayside cusp from AE-C and AE-D, J. Geophys. Res., 82, 4765, 1977. (e)
- Prangé, R., Energetic (keV) ions of ionospheric origin in the magnetosphere. A review, Ann. Géophys., 34, 187, 1978. (b,e,f)
- Prangé, R., Experimental study of the distribution function of low-energy ions (150 eV - 33 keV) in the equatorial ionosphere, during a magnetic storm associated with spread-F observations, Thesis, Doctorat d'Etat, Université Paris XI, 1978(?). (a,b)
- Price, P. B., Possible detection of energetic Jovian heavy ions at Skylab orbit, Geophys. Res. Lett., 6, 711, 1979. (e)

- Pyle, K. R., Energetic particles of planetary origin in interplanetary space, Rev. Geophys. Space Phys., 17, 587, 1979. (e)
- Pytte, T., R. L. McPherron, E. W. Hones, Jr., and H. I. West, Jr., Multiple-satellite studies of magnetospheric substorms: Distinction between polar magnetic substorms and convection-driven negative bays, J. Geophys. Res., 83, 663, 1978. (j)
- Pytte, T., R. L. McPherron, M. G. Kivelson, H. I. West, Jr., and E. W. Hones, Jr., Multiple-satellite studies of magnetospheric substorms: Plasma sheet recovery and the poleward leap of auroral zone activity, J. Geophys. Res., 83, 5256, 1978. (j)
- Pytte, T., and H. I. West, Jr., Ground-satellite correlations during presubstorm magnetic field configuration changes and plasma sheet thinning in the near-earth magnetotail, J. Geophys. Res., 83, 3791, 1978. (j)
- Raitt, W. J., R. W. Schunk, and P. M. Banks, Quantitative calculations of helium ion escape fluxes from the polar ionospheres, J. Geophys. Res., 83, 5617, 1978. (e)
- Rao, B. C. N., M. N. M. Rao, V. C. Jain, and A. Bochev, Monte-Carlo calculation of the precipitation of keV range protons and neutral hydrogen atoms into the equatorial atmosphere, Ann. Geophys., 35, 91, 1979. (a, b, h)
- Reeve, C., U. Fablesen, B. Maehlum, A. Pedersen, and P. Rothwell, Rocket measurements of ULF electric fields — and precipitating particles in the aurora, Planet. Space Sci., 26, 833, 1978. (j)
- Reiff, P. H., T. W. Hill, and J. L. Burch, Reply (to comment of W. J. Heikkila), J. Geophys. Res., 83, 229, 1978. (j)
- Richter, A. K., E. Keppler, W. I. Axford, and K. U. Denskat, Dynamics of low-energy electrons ( $\leq 17$  keV) and ions ( $\leq 80$  keV) in the vicinity of the low-latitude, duskside magnetopause: Helios 1 and 2 observations, J. Geophys. Res., 84, 1453, 1979. (b, c)
- Roelof, E. C., Solar energetic particles: From the corona to the magnetotail, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 220-241, Am. Geophys. Union, Washington, D.C., 1979. (g)
- Rönnmark, K., H. Borg, P. J. Christiansen, M. P. Gough, and D. Jones, Banded electron cyclotron harmonic instability — A first comparison of theory and experiment, Space Sci. Rev., 22, 401, 1978. (d)
- Rosenberg, T. J., K. Marthinsen, J. A. Holtet, A. Egeland, and D. L. Carpenter, Evidence of the common origin of electron microbursts and VLF chorus, J. Geomag. Geoelectr., 30, 355, 1978. (c, d)



- Rossberg, L., Undisturbed trapping boundary for energetic electrons at low altitudes, J. Geophys. Res., 83, 4307, 1978. (c)
- Rossberg, L., E. Kirsch, and M. Scholer, Intense electron and proton fluxes poleward of the trapping boundary during a substorm expansion, J. Geophys. Res., 82, 2854, 1977. (b,c)
- Rostoker, G., J. D. Winningham, K. Kawasaki, J. R. Burrows, and T. J. Hughes, Energetic particle precipitation into the high-latitude ionosphere and the auroral electrojets, 2, Eastward electrojet and field-aligned current flow at the dusk meridian, J. Geophys. Res., 84, 2006, 1979. (j)
- Roux, A., and R. Pellat, A theory of triggered emissions, J. Geophys. Res., 83, 1433, 1978. (d,h)
- Røyrvik, O., Instabilities in pitch angle diffusion and their possible relation to a 3 Hz modulation in pulsating aurora, J. Atmos. Terr. Phys., 40, 1309, 1978. (d,h)
- Ryan, J. M., M. C. Jennings, M. D. Radwin, A. D. Zych, and R. S. White, Atmospheric gamma ray angle and energy distributions from sea level to 3.5 g/cm<sup>2</sup> and 2 to 25 MeV, J. Geophys. Res., 84, 5279, 1979. (j)
- Samokhin, M. V., Generation of an Alfvén wave in the magnetotail by means of a proton accelerator, Geomagn. Aeron., 17, 469(E), 1977. (h)
- Samokhin, M. V., Possibility of producing an artificial aurora in a region that is distant in longitude from the electron beam injection site, Geomagn. Aeron., 18, 197(E), 1978. (i)
- Sari, J. W., Diurnal variation of cosmic ray intensity, 2, Calculation of the diffusion coefficients, J. Geophys. Res., 83, 5151, 1978. (g)
- Sari, J. W., D. Venkatesan, L. J. Lanzerotti, and C. G. MacLennan, Diurnal variation of cosmic ray intensity, 1, Two approaches to the study, J. Geophys. Res., 83, 5139, 1978. (g)
- Sarris, E. T., S. M. Krimigis, C. O. Bostrom, and T. P. Armstrong, Simultaneous multispacecraft observations of energetic proton bursts inside and outside the magnetosphere, J. Geophys. Res., 83, 4289, 1978. (e,g)
- Sarris, E. T., D. J. Williams, and S. M. Krimigis, Observations of counterstreaming between plasma and energetic particles in the magnetotail, J. Geophys. Res., 83, 5655, 1978. (j)
- Sato, T., Auroral and equatorial two-stream irregularities: Difference in nonlinear state, J. Geophys. Res., 82, 5195, 1977. (h)

- Scholer, M., G. Gloeckler, F. M. Ipavich, D. Hovestadt, and B. Klecker, Pitch angle distributions of energetic protons near the earth's bow shock, Geophys. Res. Lett., 6, 707, 1979. (e)
- Scholer, M., D. Hovestadt, G. Hartmann, J. B. Blake, J. F. Fennell, and G. Gloeckler, Low-altitude measurements of precipitating protons, alpha particles, and heavy ions during the geomagnetic storm on March 26-27, 1976, J. Geophys. Res., 84, 79, 1979. (b,f)
- Scholer, M., D. Hovestadt, B. Klecker, G. Gloeckler, and C. Y. Fan, Temporal development of the energetic particle composition during solar flares, J. Geophys. Res., 83, 3349, 1978. (f,g)
- Sckopke, N., and G. Paschmann, The plasma mantle: A survey of magnetotail boundary observations, J. Atmos. Terr. Phys., 40, 261, 1978. (e)
- Sentman, D. D., L. A. Frank, C. F. Kennel, D. A. Gurnett, and W. S. Kurth, Electron distribution functions associated with electrostatic emissions in the dayside magnetosphere, Geophys. Res. Lett., 6, 781, 1979. (c,d)
- Sharp, R. D., R. G. Johnson, and E. G. Shelley, Energetic particle measurements from within ionospheric structures responsible for auroral acceleration processes, J. Geophys. Res., 84, 480, 1979. (j)
- Sharp, W. E., M. H. Rees, and A. I. Stewart, Coordinated rocket and satellite measurements of an auroral event, 2, The rocket observations and analysis, J. Geophys. Res., 84, 1977, 1979. (j)
- Shavrin, P. I., To the definition of directed rigidity of geomagnetic cut-off along ASE trajectory, Geomagn. Aeron., 18, 730(R), 1978. (g)
- Shelley, E. G., Heavy ions in the magnetosphere, Space Sci. Rev., 23, 465, 1979. (f)
- Shumilov, O. I., and A. V. Frank-Kamenetskiy, Relation between the intensity of the electron flux at a geostationary orbit and the form of auroral absorption bays, Geomagn. Aeron., 17, 374(E), 1977. (c)
- Simonich, D. M., B. R. Clemesha, and W. J. H. Kirchhoff, Measurement of the north-south asymmetry in the solar proton albedo neutron flux, J. Geophys. Res., 84, 1543, 1979. (g)
- Siscoe, G. L., A quasi-self-consistent axially symmetric model for the growth of a ring current through earthward motion from a pre-storm configuration, Planet. Space Sci., 27, 285, 1979. (h)
- Smith, P. H., and N. K. Bewtra, Charge exchange lifetimes for ring current ions, Space Sci. Rev., 22, 301, 1978. (b)
- Smith, P. H., N. K. Bewtra, and R. A. Hoffman, Motions of charged particles in the magnetosphere under the influence of a time-varying large scale convection electric field, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 513-535, Am. Geophys. Union, Washington, D. C., 1979. (e)

- Solomon, J., and R. Pellat, Convection and wave-particle interactions, J. Atmos. Terr. Phys., 40, 373, 1978. (d, e, h)
- Southwood, D. J., and S. M. Kaye, Drift boundary approximations in simple magnetospheric convection models, J. Geophys. Res., 84, 5773, 1979. (e, h)
- Southwood, D. J., and R. A. Wolf, An assessment of the role of precipitation in magnetospheric convection, J. Geophys. Res., 83, 5227, 1978. (j)
- Spjeldvik, W. N., Reply (to comment of M. Walt and G. T. Davidson), J. Geophys. Res., 83, 226, 1978. (a, c)
- Spjeldvik, W. N., Expected charge states of energetic ions in the magnetosphere, Space Sci. Rev., 23, 499, 1979. (f, h)
- Spjeldvik, W. N., and T. A. Fritz, Energetic ionized helium in the quiet time radiation belts: Theory and comparison with observation, J. Geophys. Res., 83, 654, 1978. (f, h)
- Spjeldvik, W. N., and T. A. Fritz, Theory for charge states of energetic oxygen ions in the earth's radiation belts, J. Geophys. Res., 83, 1583, 1978. (f, h)
- Spjeldvik, W. N., and T. A. Fritz, Quiet time observations of equatorially trapped megaelectronvolt radiation belt ions with nuclear charge  $Z \geq 4$ , J. Geophys. Res., 83, 4401, 1978. (f)
- Stenbaek-Nielsen, H. C., and T. J. Hallinan, Pulsating auroras: Evidence of noncollisional thermalization of precipitating electrons, J. Geophys. Res., 84, 3257, 1979. (c, d)
- Stenzel, R. L., Observation of beam-generated VLF hiss in a large laboratory plasma, J. Geophys. Res., 82, 4805, 1977. (d)
- Stern, D. P., Adiabatic particle motion in a nearly drift-free magnetic field: Application to the geomagnetic tail, J. Geophys. Res., 83, 1079, 1978. (e, h)
- Swift, D. W., Turbulent generation of electrostatic fields in the magnetosphere, J. Geophys. Res., 82, 5143, 1977. (d, h)
- Swift, D. W., Acceleration mechanisms for auroral electrons, J. Geomag. Geoelectr., 30, 449, 1978. (h)
- Swift, D. W., Auroral mechanisms and morphology, Rev. Geophys. Space Phys., 17, 681, 1979. (j)
- Szuszcwicz, E. P., D. N. Walker, J. C. Holmes, and H. Leinbach, Plasma diffusion in a space-simulation beam-plasma-discharge, Geophys. Res. Lett., 6, 201, 1979. (d, i)

- Tamao, T., Modulation of energetic particle fluxes due to long-period geomagnetic oscillations, Planet. Space Sci., 26, 1149, 1978. (c)
- Thorne, R. M., S. R. Church, and D. J. Gorney, On the origin of plasmaspheric hiss: The importance of wave propagation and the plasma-pause, J. Geophys. Res., 84, 5241, 1979. (d,h)
- Thorne, R. M., and B. T. Tsurutani, Power-line harmonic radiation: Can it significantly affect the earth's radiation belts?, Science, 204, 839, 1979. (i)
- Tinsley, B. A., Energetic neutral atom precipitation during magnetic storms: Optical emission, ionization, and energy deposition at low and middle latitudes, J. Geophys. Res., 84, 1855, 1979. (b,f)
- Tolba, M. F., On the latitude dependence of the semidiurnal anisotropy of cosmic rays, Planet. Space Sci., 27, 355, 1979. (g)
- Topchygin, I. N., Acceleration of particles by shocks in a cosmic plasma, Space Sci. Rev., in press, 1980. (j)
- Torbert, R. B., and F. S. Mozer, Electrostatic shocks as the source of discrete auroral arcs, Geophys. Res. Lett., 5, 135, 1978. (j)
- Torr, M. R., and D. G. Torr, Energetic oxygen: A direct coupling mechanism between the magnetosphere and thermosphere, Geophys. Res. Lett., 6, 700, 1979. (f)
- Trefall, H., and D. J. Williams, Time structure of postmidnight energetic electron precipitation and the limit of stable trapping, J. Geophys. Res., 84, 2725, 1979. (c,d)
- Tsurutani, B. T., S. R. Church, and R. M. Thorne, A search for geographic control of the occurrence of magnetospheric ELF emissions, J. Geophys. Res., 84, 4116, 1979. (i)
- Isutsui, M., K. Hiramoto, H. Matsumoto, and I. Kimura, VLF plasma wave excitation due to low-energy electron beam injection: Laboratory simulation of a rocket experiment, J. Geophys. Res., 84, 4217, 1979. (d,i)
- Tulinov, V. F., V. M. Feygin, M. A. Savel'yev, Yu. M. Zhuchenko, V. A. Lipovetskiy, V. A. D'yachenko, A. P. Babayev, T. A. Zhuchenko, V. V. Tulyakov, and L. S. Novikov, Experimental investigation of the daily variations of precipitating corpuscular emissions, Geomagn. Aeron., 17, 333(E), 1977. (j)
- Ungstrup, E., D. M. Klumpar, and W. J. Heikkila, Heating of ions to superthermal energies in the topside ionosphere by electrostatic ion cyclotron waves, J. Geophys. Res., 84, 4289, 1979. (d,h)

- Vampola, A. L., VLF transmission induced slot electron precipitation, Geophys. Res. Lett., 4, 569, 1977. (ii)
- Vampola, A. L., and G. A. Kuck, Induced precipitation of inner zone electrons, 1. Observations, J. Geophys. Res., 83, 2543, 1978. (a,c)
- Velinov, P., Profiles of ionization of low-energy solar and magnetospheric particles in the ionosphere, Geomagn. Aeron., 18, 31(E), 1978. (g)
- Venkatarangan, P., and D. I. McEwen, Electron measurements (18 keV - 20 eV) in auroral events, Planet. Space Sci., 27, 669, 1979. (cc)
- Veselovskiy, I. S., Movement of a beam of charged particles in the earth's magnetosphere, Geomagn. Aeron., 17, 326(E), 1977. (ee)
- Vette, J. I., Overview — High energy particles, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 118-129. Am. Geophys. Union, Washington, D. C., 1979. (ji)
- Viswanathan, K. S., and G. Renuka, Pitch angle diffusion by bounce resonance, Planet. Space Sci., 26, 75, 1978. (d,h)
- Volodichev, N. N., E. I. Morozova, M. A. Saraeva, A. A. Suslov, and P. I. Shavrin, To the definition of the spectrum of the proton flare of August 4, 1972, by the latitudinal effect on artificial earth satellite Prognoz-2, Geomagn. Aeron., 19, 732-R, 1979. (g)
- Volokitin, A. S., and E. V. Mashin, About relaxation of an intense ray of energetic electrons in the ionosphere, Geomagn. Aeron., 19, 739(R), 1979. (j)
- Voss, H. D., and L. G. Smith, Nighttime ionization by energetic particles at Wallops Island in the altitude region 120 to 260 km, Geophys. Res. Lett., 6, 93, 1979. (j)
- Wagner, J. S., J. R. Kan, and S.-I. Matsufu, Particle dynamics in the plasma sheet, J. Geophys. Res., 84, 891, 1979. (e,h)
- Walker, R. J., K. N. Erickson, and J. R. Winckler, Pitch angle dispersion of drifting energetic protons at synchronous orbit, J. Geophys. Res., 83, 1595, 1978. (b,e)
- Wallis, D. D., J. R. Burrows, M. C. Mushup, C. D. Anger, and J. S. Murphree, Observations of particles precipitating into detached arcs and patches equatorward of the auroral oval, J. Geophys. Res., 84, 1347, 1979. (c)
- Walt, M., and G. T. Davidson, Comment on "Radiation belt electrons: Structure of the loss cone" by W. N. Spiedvik, J. Geophys. Res., 83, 225, 1978. (a,c)
- Walt, M., L. L. Newkirk, and W. E. Francis, Bremsstrahlung produced by precipitating electrons, J. Geophys. Res., 84, 967, 1979. (cc)

- Weber, E. J., J. A. Whalen, R. A. Wagner, and J. Buchau, A 12-hour case study of auroral phenomena in the midnight sector: Electrojet and precipitating particle characteristics, J. Geophys. Res., 82, 3557, 1977. (j)
- West, H. I., Jr., The signatures of the various regions of the outer magnetosphere in the pitch angle distributions of energetic particles, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 150-179, Am. Geophys. Union, Washington, D.C., 1979. (c)
- West, H. I., Jr., R. M. Buck, and M. G. Kivelson, On the configuration of the magnetotail near midnight during quiet and weakly disturbed periods: State of the magnetosphere, J. Geophys. Res., 83, 3805, 1978. (j)
- West, H. I., Jr., R. M. Buck, and M. G. Kivelson, On the configuration of the magnetotail near midnight during quiet and weakly disturbed periods: Magnetic field modeling, J. Geophys. Res., 83, 3819, 1978. (j)
- Whalen, B. A., W. Bernstein, and P. W. Daly, Low altitude acceleration of ionospheric ions, Geophys. Res. Lett., 5, 55, 1978. (e)
- Whalen, B. A., and P. W. Daly, Do field-aligned auroral particle distributions imply acceleration by quasi-static parallel electric fields?, J. Geophys. Res., 84, 4175, 1979. (j)
- Whipple, E. C., Jr., (U, B, K) coordinates: A natural system for studying magnetospheric convection, J. Geophys. Res., 83, 4318, 1978. (e,h)
- Whipple, E. C., Jr., and S. E. DeForest, Overview — Low energy particles, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 338-339, Am. Geophys. Union, Washington, D.C., 1979. (j)
- Whipple, E. C., Jr., A kinetic approach to magnetospheric modeling, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 462-472, Am. Geophys. Union, Washington, D.C., 1979. (e,h)
- Wilken, B., T. A. Fritz, A. Korth, and G. Kremser, Substorm-associated injections of energetic ions observed by GEOS-1 and -2 in and near synchronous altitude, Space Sci. Rev., 22, 647, 1978. (a)
- Williams, D. J., Magnetopause characteristics inferred from three-dimensional energetic particle distributions, J. Geophys. Res., 84, 101, 1979. (b,c)
- Williams, D. J., Observations of significant magnetosheath antisolar energy flow, J. Geophys. Res., 84, 2105, 1979. (j)

- Williams, D. J., and L. A. Frank, ISEE 1 charged particle observations indicative of open magnetic field lines near the subsolar region, NOAA/SEL preprint, Boulder, Colo., September 1979. (j)
- Williams, D. J., T. A. Fritz, E. Keppler, B. Wilken, and G. Wibberenz, Three-dimensional energetic ion observations and the tangential magnetopause electric field, Space Sci. Rev., 22, 805, 1978. (b)
- Williams, D. J., T. A. Fritz, B. Wilken, and E. Keppler, An energetic particle perspective of the magnetopause, NOAA/SEL preprint, Boulder, Colo., March 1979. (j)
- Winningham, J. D., C. D. Anger, G. G. Shepherd, E. J. Weber, and R. A. Wagner, A case study of the aurora, high-latitude ionosphere, and particle precipitation during near-steady state conditions, J. Geophys. Res., 83, 5717, 1978. (j)
- Winningham, J. D., K. Kawasaki, and G. Rostoker, Energetic particle precipitation into the high-latitude ionosphere and the auroral electrojets, 1, Definition of electrojet boundaries using energetic electron spectra and ground-based magnetometer data, J. Geophys. Res., 84, 1993, 1979. (c)
- Yemel'yanenko, S. N., S. N. Kuznetsov, G. B. Lopatina, and V. G. Stolpovskiy, Fast pulsations of electron density in the trapping region, Geomagn. Aeron., 18, 348(E), 1978. (c)
- Yemel'yanenko, S. N., S. N. Kuznetsov, and V. G. Stolpovskiy, Outer radiation belt during a strong magnetic storm, Geomagn. Aeron., 18, 141(E), 1978. (c)
- Young, D. T., Ion composition measurements in magnetospheric modeling, in Quantitative Modeling of Magnetospheric Processes, edited by W. P. Olson, pp. 340-363, Am. Geophys. Union, Washington, D.C., 1979. (b,f)
- Young, D. T., J. Geiss, H. Balsiger, P. Eberhardt, A. Ghielmetti, and H. Rosenbauer, Discovery of  $\text{He}^{2+}$  and  $\text{O}^{2+}$  ions of terrestrial origin in the outer magnetosphere, Geophys. Res. Lett., 4, 561, 1977. (f)
- Zakharchenko, V. F., and Yu. Ya. Polyak, Influence of anisotropy of a magnetospheric plasma on long-period pulsations, Geomagn. Aeron., 19, 111(R), 1979. (b)
- Zakharov, A. V., and S. N. Kuznetsov, Lifetime of the earth's radiation belt electrons and VLF emission, Geomagn. Aeron., 18, 232(E), 1978. (c,d)
- Zuyeva, Z. G., V. I. Lazareva, and M. V. Fel'tsov, Location of the neck of the auroral electron discontinuity in the morning sector of the magnetosphere according to measurements on the Molniya-1 satellite, Geomagn. Aeron., 17, 622(E), 1977. (c)

Zwickl, R. D., E. C. Roelof, R. E. Gold, S. M. Krimigis, and  
I. P. Armstrong, Z-rich solar particle event characteristics  
1972-1976, Astrophys. J., 225, 281, 1978. (f, g)



## LABORATORY OPERATIONS

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the Nation's rapidly developing space systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

Aerophysics Laboratory: Aerodynamics; fluid dynamics; plasmadynamics; chemical kinetics; engineering mechanics; flight dynamics; heat transfer; high-power gas lasers, continuous and pulsed, IR, visible, UV; laser physics; laser resonator optics; laser effects and countermeasures.

Chemistry and Physics Laboratory: Atmospheric reactions and optical backgrounds; radiative transfer and atmospheric transmission; thermal and state-specific reaction rates in rocket plumes; chemical thermodynamics and propulsion chemistry; laser isotope separation; chemistry and physics of particles; space environmental and contamination effects on spacecraft materials; lubrication; surface chemistry of insulators and conductors; cathode materials; sensor materials and sensor optics; applied laser spectroscopy; atomic frequency standards; pollution and toxic materials monitoring.

Electronics Research Laboratory: Electromagnetic theory and propagation phenomena; microwave and semiconductor devices and integrated circuits; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, superconducting and electronic device physics; millimeter-wave and far-infrared technology.

Materials Sciences Laboratory: Development of new materials; composite materials; graphite and ceramics; polymeric materials; weapons effects and hardened materials; materials for electronic devices; dimensionally stable materials; chemical and structural analyses; stress corrosion; fatigue of metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere; the effects of optical, electromagnetic, and particulate radiations in space on space systems.

DATE  
FILMED  
1987